

1 **Minamata Convention on Mercury**
2
3 **DRAFT Report on the work of the ad hoc technical group on effectiveness**
4 **evaluation**
5
6 **Open for comment: 1 August to 5 September 2019**
7 Comments to: mea-minamatasecretariat@un.org
8
9 Enquiries to be sent to: Claudia.tenHave@un.org and Eisaku.Toda@un.org
10
11

Table of Content	
Executive summary	
I Introduction	
II Overall description of the effectiveness evaluation framework	
III Proposed methodology and schedule for the evaluation	
IV Issues for further consideration	
Annex 1: Technical information on monitoring	
Annex 2: Draft terms of reference of the effectiveness evaluation committee	
Annex 3: Draft terms of reference of the global monitoring arrangements	
Annex 4: Description of the reports to be prepared for the Effectiveness Evaluation Committee	

12
13

14 **Executive Summary**

15
16 **Introduction**

17
18 This report proposes a framework for the effectiveness evaluation and monitoring
19 arrangements under the Minamata Convention on Mercury. The report was developed by the
20 ad hoc technical expert group based on mandates provided in decisions MC-1/9 and MC-
21 2/10, and information contained in submissions by Parties, stakeholders and other
22 information. The report represents the outcome of consultations and review performed by
23 experts, including two meetings of the ad hoc group in 2018 and 2019 respectively, with
24 follow-up drafting and reviewing by experts and commentators.

25
26 Building on the identification of elements for the effectiveness evaluation framework and
27 review of monitoring activities included in the first report from the ad hoc group to COP-2,
28 this report proposes a framework setting out arrangements, information flows, and the
29 required reports on which the Effectiveness Evaluation Committee will base its consideration
30 of the effectiveness of the Convention for presentation to the Conference of the Parties.

31
32 **Effectiveness evaluation framework**

33
34 In sections II and III, the report elaborates on the proposed organization of the effectiveness
35 evaluation framework.

36
37 The effectiveness evaluation is based on four policy questions that will allow the Conference
38 of the Parties to consider whether the Convention will achieve its objective of protecting
39 human health and the environment from the anthropogenic emissions and releases of mercury
40 and mercury compounds. The policy questions are:

- 41 1. Have the Parties taken actions to implement the Minamata Convention?
- 42 2. Have the actions resulted in changes in emission and releases of mercury to the
43 environment?
- 44 3. Have these changes in emission and releases resulted in changes in levels of
45 mercury in the environment, biota and humans attributable to the Convention?
- 46 4. Will existing measures under the Minamata Convention be sufficient to meet its
47 objectives of promoting human health and environment from mercury?

48
49 The framework relies on evaluating evidence along the causal pathway linking actions to
50 implement the Convention, associated changes in emissions and releases, and resulting
51 changes in levels and trends in the global environment, biota, and humans. The ad hoc group
52 proposes sets of indicators on process, outcome and monitoring to inform these policy
53 questions. The proposed indicators draw on previous work on elements of the effectiveness
54 evaluation framework and the review of monitoring capacities and abilities.

55
56 Numerous indicators, developed following an article-by-article review, are clustered to
57 enable synthesized analysis for an integrated picture. Specific articles of central or
58 crosscutting importance to the overall provisions are not clustered (such as Art 1 and 16). The
59 following articles and indicators are clustered:

- 60 ▪ **Supply cluster:** supply, storage and waste (Art 3, 10, 11);
- 61 ▪ **Demand cluster:** products, processes and ASGM (Art 4,5,7);
- 62 ▪ **Pressures cluster:** emissions, releases and contaminated sites (Art 8, 9, and 12),
- 63 ▪ **Support cluster:** financial and technical assistance (Art 13 and 14), and
- 64 ▪ **Information and research cluster:** information exchange, public information,
- 65 research (Art 17, 18 and 19).

66 Article 22, paragraph 3 indicates that the evaluation shall be conducted using available
67 scientific, environmental, technical, financial and economic information. Two streams of
68 information are referred to in this regard: (i) information provided by Parties based on Article
69 21 reporting, and (ii) information and knowledge that is scientific, peer-reviewed and
70 publicly available.
71

72
73 Based on such information the framework foresees five reports to be produced (see
74 description in Section III and Annex 4). The following four synthesis reports are to be
75 prepared:

- 76 1. The state of global mercury levels in the environment, biota and humans, as well
77 as trends over time, that is a **Global Monitoring Report**;
- 78 2. **Emissions and Releases** – modelled after the *Global Mercury Assessment* (2018)
- 79 3. **Trade, Supply and Demand** – modelled after the report on *Global Mercury:*
80 *Supply, Trade and Demand* (2017);
- 81 4. **Waste Management** – building on the *Global Mercury Waste Assessment* (2017).

Deleted: 8

82
83 The framework presents a flow of information from level 1 to level 6, namely starting from
84 collecting information, to compiling information, to analysing and synthesizing information
85 (levels 1 to 3). The framework then foresees an integrating function (level 4) before reaching
86 the Effectiveness Evaluation Committee (level 5) and the Conference of the Parties (level 6).

87
88 The framework identifies different entities that have different roles in the process. While
89 some entities already exist (i.e. those for administrative and programme support, compilation
90 of data for synthesis reports, etc.) there are others who are to perform vital scientific and
91 technical analyses to implement the framework, but they are not in place yet.

92
93 The framework foresees two scientific and technical functions to perform analysis, synthesis
94 and interpretation at levels 3 and 4: The first function (level 3) is to synthesise mercury
95 information collected and compiled. This function foresees a role for the secretariat, for
96 scientists and experts, and for organisations. The second function (level 4) is to interpret the
97 information and knowledge collected and synthesised, to interpret the linkages between
98 policy actions, emission reductions and resulting mercury levels, using available data sources,
99 modelling techniques and analytical tools drawn from natural and social sciences. The
100 function foresees the production of the **Integrated Assessment Report** for the Effectiveness
101 Evaluation Committee. This report is to be science-based but should be accessible to non-
102 technical readers.

103
104 The Effectiveness Evaluation Committee (see its Terms of Reference in Annex II of the
105 report) will consider all the reports produced to consider the policy questions outlined above

107 and derive conclusions about the effectiveness of the Convention for its report to the
108 Conference of the Parties.

109
110 Thereafter, the Conference of the Parties receives and reviews the report of the Effectiveness
111 Evaluation Committee, and considers the conclusions and recommendations for the
112 Convention.

113
114 The framework is submitted to the third meeting of the Conference of the Parties for
115 adoption, and foresees a timeline for the first cycle of the effectiveness evaluation where the
116 fourth meeting of the Conference of the Parties establishes the Effectiveness Evaluation
117 Committee, and the fifth meeting of the Conference of the Parties considers the findings of
118 that Committee.

119
120 The report also outlines in Section IV further issues for the Conference of the Parties to
121 consider at its third meeting, including a proposed draft decision.

122 123 **Monitoring arrangements**

124
125 The report addresses the task outlined in decisions MC-1/9 and MC-2/10 in relation to
126 monitoring, by reviewing available monitoring data, identifying gaps, examining modelling
127 capabilities, and outlining global monitoring arrangements.

128
129 In considering monitoring information data, the ad hoc group considered matrices mentioned
130 in MC-2/10: air, humans, biota and water. The ad hoc group concluded that data on levels of
131 mercury in air, biota and humans either are available or would be able to be obtained and
132 comparable on a global basis.

133
134 Levels of mercury and mercury compounds in water are collected in relation to water quality
135 issues in a number of countries. These data may be useful in tracking mercury resulting from
136 local activities which release mercury; however, it will not provide overall trends on a global
137 basis. Levels of mercury in ocean water could be comparable on a global basis and collected
138 by existing networks and ad hoc research programmes, but currently such work is done
139 through research-based activities and not dedicated long-term monitoring programmes

140
141 The global modelling capabilities have been reviewed in order to understand availability of
142 tools for the use in the effectiveness evaluation. The models complement monitoring data
143 with estimation based on scientific understanding of mechanisms affecting mercury
144 behaviour. Models for different media (air, water, land, biota) vary in their ability and state of
145 development. Atmospheric models have been extensively evaluated and can be applied to
146 assess spatial gradients in atmospheric mercury concentrations and deposition, as well as
147 temporal changes. By contrast, models for other media such as land are still mainly used in
148 research applications. Integrated models that work across media drawing on expertise that
149 bridges natural science, social science, and engineering, are undergoing rapid development in
150 the scientific and academic community and are expected to be available by 2023 for policy-
151 relevant analyses.

152

153 In the consideration of the monitoring arrangements, the following key elements were
154 identified:

- 155 ▪ Mercury data and their availability from human health and environmental
156 monitoring programmes that achieve global coverage and contain at least core
157 representative data from all regions,
- 158 ▪ Tools supporting data harmonization such as standard operating procedures
159 and monitoring guidance document,
- 160 ▪ Expertise necessary for gathering and consolidating harmonized information
161 that ensures comparability and consistency in monitoring data over the long-
162 term,
- 163 ▪ Modelling capabilities, and
- 164 ▪ Development of a global periodic report to support the effectiveness
165 evaluation.

166
167 In line with the proposal to perform scientific functions and to carry out tasks related to work
168 with monitoring indicators identified in the effectiveness evaluation framework, scientific
169 expertise and qualifications are required to oversee the gathering and consolidation of
170 monitoring data. It is proposed that this task should be overseen by an expert body whose
171 terms of reference are proposed in Annex 3.

172
173 **NOTE:** In addition to this report, the ad hoc group developed a complementing information
174 document. Part 1 of that document provides a more detailed review of available monitoring
175 data and background on the proposal for monitoring activities with further scientific and
176 technical details. Part 2 of the information document contains a proposal for elements of the
177 guidance document for mercury monitoring that will be developed under the monitoring
178 arrangements to be established by the Conference of the Parties.

179
180

181 **I. Introduction**

182

183 1. At the first and second meetings, the Conference of the Parties tasked an ad hoc
184 expert group to consider the arrangements to be put in place to provide the
185 Conference of the Parties with the required information to conduct an effectiveness
186 evaluation of the Minamata Convention on Mercury.¹ The effectiveness evaluation is
187 to be conducted at regular intervals, with the first taking place within six years after
188 the entry into force of the Convention.² This report is the outcome of the consultation
189 and deliberations of the ad hoc expert group to put in place arrangements to conduct
190 an effectiveness evaluation.

191 2. Article 22 of the Minamata Convention in paragraph 2 stipulates that the Conference
192 of the Parties, shall initiate the establishment of arrangements for providing itself with
193 comparable monitoring data on the presence and movement of mercury and mercury
194 compounds in the environment, as well as the trends in the levels of mercury and
195 mercury compounds as observed in biotic media and vulnerable populations.
196 Paragraph 3 of that article further stipulates that the evaluation shall be conducted
197 based on available scientific, environmental, technical, financial and economic
198 information, including:

199 (a) Reports and other monitoring information provided to the Conference of
200 the Parties pursuant to paragraph 2;

201 (b) Reports submitted pursuant to Article 21;

202 (c) Information and recommendations provided pursuant to Article 15; and

203 (d) Reports and other relevant information on the operation of the financial
204 assistance, technology transfer and capacity building arrangements put in place
205 under this Convention.

206 3. The first meeting of the Conference of the Parties recognised the urgent need for a
207 framework for the effectiveness evaluation that includes a strategic, cost-effective
208 approach that provides appropriate and sufficient data, and further acknowledged
209 publications such as UNEP's global mercury assessments, as well as the GEF-funded
210 Minamata Initial Assessments, as important sources of information. The Conference
211 of the Parties set out a roadmap which included the establishment of the ad hoc group

¹ MC-1/9 on the Establishment of arrangements in regard to effectiveness evaluation established the ad hoc group of experts on the arrangements for providing the Conference of the Parties with comparable monitoring data, and elements of an effectiveness evaluation framework under article 22 of the Minamata Convention (hereafter referred to as the ad hoc expert group). The decision also laid out a roadmap for establishing arrangements for providing the Conference of the Parties with comparable monitoring data, and elements of an effectiveness evaluation framework under article 22 of the Minamata Convention. The ad hoc expert group produced a report which was presented to the second meeting of the Conference of the Parties (See UNEP/MC/COP.2/13 and UNEP/MC/COP.INF/8).

Subsequently, MC-2/10 extended the terms of reference of the ad hoc technical expert group, adopted a roadmap for the subsequent work, requested the ad hoc expert group to develop the terms of reference for global monitoring arrangements, and requested the ad hoc expert group to report its progress to the third meeting of the Conference of the Parties.

² While the first effectiveness evaluation of the Minamata Convention on Mercury is to take place within six years of the entry into force of the Convention, the Conference of the Parties is to decide on the future interval of the effectiveness evaluations.

212 of experts on the arrangement for providing the Conference of the Parties with
 213 comparable monitoring data, and the elements of an effectiveness evaluation
 214 framework (see MC-1/9 Establishment of arrangement in regard to effectiveness
 215 evaluation).

216 4. For deliberation of these matters and based on the roadmap and terms of reference
 217 outlined in MC-1/9, the ad hoc group of experts began its work at its first meeting in
 218 Ottawa, Canada (5-9 March 2018). The outcome of the work of this first round of
 219 deliberations, reflecting comments received during the subsequent open comment
 220 period, was presented to the second meeting of the Conference of the Parties in
 221 Geneva in November 2018 (see UNEP/MC/COP.2/13 and UNEP/MC/COP.2/INF/8).

222 5. The second meeting of the Conference of Parties deliberated on the outcome of the ad
 223 hoc group of experts and decided to revise the Group's mandate and identify
 224 additional expertise needed to enable it to complete its work for presentation to the
 225 third meeting of the Conference of the Parties in November 2019. The Conference of
 226 the Parties in its decision 2/10 also requested the ad hoc expert group to undertake the
 227 following tasks:

228 (a) Using the objective of the Minamata Convention, review and assess the
 229 detailed article-by-article process and outcome indicators presented in
 230 UNEP/MC/COP.2/INF/8, and elaborate on the sources of information and
 231 baselines for those indicators, considering cost-effectiveness, practicality,
 232 feasibility and sustainability, and, on that basis, provide detailed rationales for
 233 the recommended indicators;

234 (b) Identify which recommended indicators require monitoring data, in particular
 235 in relation to the control measures and objectives set out in the articles of the
 236 Convention;

237 (c) Develop a methodology for integrating the recommended indicators with a
 238 view to providing an integrative picture of the general effectiveness of the
 239 Convention, (e.g., by use of cross-cutting indicators); and

240 (d) Amend the recommended draft terms of reference of the effectiveness
 241 evaluation committee and the schedule for the first effectiveness evaluation, if
 242 needed, on the basis of the outcome of the above.

243 6. Following its revised mandate, the re-named ad hoc technical working group met in
 244 Geneva in April 2019 to deliberate specifically on the requested report to be presented
 245 to the third meeting of the Conference of the Parties. The present report is the
 246 outcome of the work begun at that meeting³ and completed in the subsequent months
 247 that included an open comment period from 1 August to 5 September 2019.

248 7. Following the guidance of MC-2/10, this report is presented in four sections: Section I
 249 gives an introduction on the mandate of the work of the ad hoc technical expert group,
 250 and the report on its work on the arrangements the group proposes be put in place to
 251 provide the Conference of the Parties with the required information to conduct an

³ At this meeting, the ad hoc technical group had before it a compilation of comments on the effectiveness evaluation framework (UNEP/MC/EE.2/5), submitted information (UNEP/MC/EE.2/3) and the compilation of comments on the report of the group submitted to the Conference of the Parties at its second session (UNEP/MC/EE.2/4).

252 effectiveness evaluation of the Minamata Convention on Mercury. Section II provides
253 an overview description of the proposed effectiveness evaluation framework
254 including laying out four key policy question identified to be used to evaluate the
255 effectiveness of the Convention. Section III lays out the constituent elements of the
256 framework in detail by describing the proposed methodology to conduct the
257 effectiveness evaluation. This section also puts forward the proposed schedule.
258 Section IV outlines further issues for the consideration by the Conference of the
259 Parties, including as suggested action a draft decision to operationalise the proposed
260 framework.

261 8. The report further contains four annexes. The first annex outlines technical
262 information related to monitoring. The second annex presents the terms of reference
263 of the Effectiveness Evaluation Committee. The third annex outlines the terms of
264 reference of the global monitoring arrangements. And the fourth annex gives a
265 description of the reports that are to be prepared for the Effectiveness Evaluation
266 Committee.

267 9. The ad hoc technical expert group proposes that the Conference of the Parties at its
268 third meeting adopts the framework, adopts the proposed timeline for the first cycle of
269 the effectiveness evaluation, and at its fourth meeting establishes the Effectiveness
270 Evaluation Committee. This will enable the fifth meeting of the Conference of the
271 Parties to consider the findings and conclusions of the Effectiveness Evaluation
272 Committee. A draft decision has been prepared for consideration in this regard.

273

274 **II. Overview description of the effectiveness evaluation framework**

275

276 10. The objective of the Minamata Convention, per Article 1, is “*to protect the human*
277 *health and the environment from anthropogenic emissions and releases of mercury*
278 *and mercury compounds.*”

279 11. The goal of an effectiveness evaluation is to consider the extent to which the
280 Convention is achieving this objective. To analyse its effectiveness is to consider
281 whether measures taken by Parties in response to the Convention have resulted in
282 reductions in emissions and releases that have, in turn, led to lower risks to human
283 health and the environment (compared with what would have occurred if the
284 Convention would not have been implemented).

285 12. The framework for the effectiveness evaluation of the Minamata Convention, as
286 proposed by the ad hoc technical expert group, relies on evaluating evidence along the
287 causal pathways linking actions to implement the Convention, associated changes in
288 emissions and releases, and resulting changes in levels and trends in the global
289 environment, biota, and humans. ⁴

290 13. Based on the information collated, and through proposed indicators on process,
291 outcome and monitoring, an assessment will be made of whether changes in mercury
292 levels attributable to the Convention are significant and sufficient in relations to four
293 policy questions.

294

295 **Policy Questions**

296 14. **First Policy Question: Have the Parties taken actions to implement the**
297 **Minamata Convention?** The framework contains a succinct set of “process”
298 indicators intended to reflect the level of implementation of the Convention by
299 Parties. These indicators can be used to evaluate whether implementation of
300 Convention measures can be credibly linked to changes in emissions and releases.
301 They can also be used to identify common challenges in implementation that may
302 undermine the Convention effectiveness. The process indicators are based primarily
303 on reporting mandated by the Convention, supplemented by other available scientific,
304 environmental, technical, financial and economic information as per Article 22,
305 paragraph 3.

306

307 15. **Second Policy Question: Have these actions resulted in changes in emissions and**
308 **releases of mercury to the environment?** The framework also contains a set of
309 “outcome” indicators that reflect estimated changes in supply, demand and emissions
310 and releases of Hg due to Convention measures, as reported by Parties under the
311 Convention. The framework suggests supplementing these data with context provided
312 by comprehensive estimates of global mercury supply, demand, emissions and
313 releases.

314

315 16. **Third Policy Question: Have these changes in emissions and releases resulted in**
316 **changes in levels of mercury in the environment, biota and humans attributable**
317 **to the Convention?** Article 22 of the Convention specifies that monitoring data on

318 “the presence and movement of mercury and mercury compounds in the environment
319 as well as trends in levels of mercury and mercury compounds observed in biotic
320 media and vulnerable populations,” should be used in the effectiveness evaluation.
321 Attributing changes in human and environmental levels of mercury to Convention
322 measures is challenging, but possible. Past and present emissions from human
323 activities combine with natural sources and other factors affecting mercury cycling,
324 such as atmospheric and ecosystem characteristics, which may evolve, inter alia, due
325 to climate change. The framework outlines how global mercury measurements can be
326 used to assist in the assessment of how successful the implementation measures of the
327 Convention have been.

328
329 17. This complex system results in the observed levels of mercury in the environment,
330 biota and humans. As scientific knowledge is still developing to better directly link
331 sources to these receptors, integrated modelling approaches are needed to estimate
332 how changes of emissions and releases from sources covered by the Convention
333 contribute to changes in levels in the environment, biota and humans. The ongoing
334 development and validation of such integrated models relies on monitoring data as
335 well as scientific knowledge of environmental processes and will assist in attributing
336 mercury changes in the environment, to change in mercury emissions and releases.
337

338 **18. Fourth Policy Question: Will existing measures under the Minamata Convention**
339 **be sufficient to meet its objectives of protecting human health and the**
340 **environment from mercury?** The response to the third policy question will tell us to
341 what extent the Convention is affecting levels and trends of mercury in the
342 environment, biota and humans. The fourth policy question will look at whether the
343 measures under the Convention is significant and sufficient. Is the Convention
344 delivering reduced emissions and releases to its full potential? If not, why? Would
345 delivering at full potential prevent the majority or only a small part of anthropogenic
346 emissions and releases of mercury? Furthermore, are management measures to
347 address residual risk adequate and sufficient in addressing the exposure of people to
348 mercury?

349

350 **Expert-led integrated assessment for consideration by the Effectiveness Evaluation**
351 **Committee**

352 19. The proposed framework envisions the synthesis of information and knowledge in
353 **synthesis reports**, and based on these reports and additional information, the
354 preparation of an **integrated assessment report** that provides a scientific and
355 technical perspective on the four policy questions articulated above. The integrated
356 assessment report will interpret the linkages between policy actions, emissions
357 reductions and resulting mercury levels, using available data sources, modelling
358 techniques, and analytical tools drawn from natural and social sciences, and other
359 relevant knowledge.

360 20. Present science has not yet developed reliable models for forecasting long-term
361 changes in mercury levels resulting from emissions reductions that take into account
362 the full complexities of mercury in the environment. Pending the availability of
363 suitable confirmed multi-media models, the integrated assessment report for the first
364 evaluation may or may not include the use of forecasting models. Therefore, earlier
365 evaluations on the effectiveness of the Minamata Convention may have greater

366 uncertainty than later evaluations when improvements to such forecasting models
367 become available.

368 21. The Effectiveness Evaluation Committee will use the expert-led scientific and
369 technical integrated assessment report, and supplemented as necessary by the other
370 synthesis reports, to consider the policy questions and to draw conclusions about the
371 effectiveness of the Convention. Based on this evaluation, the Effectiveness
372 Evaluation Committee is to make recommendations to the Conference of the Parties
373 as required.

374 22. **Table 1** below gives an overview of the construction of the overall effectiveness
375 evaluation framework from the above-mentioned four policy questions, to indicators
376 proposed for evaluation, to the required synthesis reports and the integrated
377 assessment report that will be prepared for the Effectiveness Evaluation Committee,
378 for its consideration and report to the Conference of the Parties. The constituent
379 elements of the framework are explained in detail in Section II.

380
381

Commented [m1]: Modification suggested: The table contains information inconsistent with the body text.

Table 1: Construction of the effectiveness evaluation framework from policy questions, to indicators and to required reports for consideration by the Effectiveness Evaluation Committee

Policy Questions	First Policy Question: Have the Parties taken actions to implement the Minamata Convention?	Second Policy Question: Have these actions resulted in changes in emissions and releases of mercury to the environment?	Third Policy Question: Have these changes in emissions and releases resulted in changes in levels of mercury in the environment, biota and humans attributable to the Convention?	Fourth Policy Question: Will existing measures under the Minamata Convention be sufficient to meet its objectives of protecting human health and the environment from mercury?
Indicators	Process indicators (<i>para 46</i>)	Outcome indicators (<i>para 46</i>) Monitoring indicators (<i>para 46</i>)	Monitoring indicators (<i>para 52</i>)	
Indicator Clusters	1. Supply Cluster 2. Demand Cluster 3. Pressure Cluster ---- 4. Support Cluster 5. Info and Research Cluster	1. Supply Cluster 2. Demand Cluster 3. Pressure Cluster ---- 4. Support Cluster 5. Info and Research Cluster	1. Pressure Cluster	1. independent Article 1
Information Sources	Parties: Article 21 reports (<i>main source</i>)	Parties: Article 21 reports (<i>main source</i>)	- Parties: Article 21 reports - Existing/proposed monitoring networks and models	- Academic articles and other information on socio-economic, technology, climate, global policies, etc. - Emissions and releases - Trade, supply and demand - Waste management - Monitoring report
Secretariat documents to COP, according to Article 22	- ICC reports - Financial mechanism reports - Report on Capacity-building and technical assistance	n/a	n/a	
Reports prepared for the Effectiveness Evaluation Committee	Level 1 – 3		Level 3	
	1. Emissions and Releases (Pressure Cluster) “ <i>Mercury to the environment</i> ” 2. Trade, Supply and Demand (<i>Supply and Demand Clusters</i>) “ <i>Intended/economic movement of mercury</i> ” 3. Waste Management (Supply, Demand and Pressure Clusters)		4. Global Monitoring Report	
	Level 4			
	5. Integrated assessment Report			

Formatted Table

	<p>Report of the Effectiveness Evaluation Committee is considered by the Conference of the Parties</p> <p><u>The Effectiveness Evaluation Committee will use the Integrated Assessment Report supplemented by the synthesis reports* to consider the policy questions posed in the framework, and from that derive conclusions about the effectiveness of the Convention.</u></p>
--	--

Level 5


Deleted: 6

384 **III. Proposed methodology and schedule for the evaluation**

385
386 **1. Information and analysis flow for the proposed effectiveness evaluation**

387
388 23. The effectiveness evaluation will be carried out through a series of sequential steps of
389 data identification and collection, compilation of information, assessment, analysis and
390 synthesis. The framework presents the flow of information, beginning with identifying
391 and collecting information, to compiling information, to synthesises information (levels 1
392 to 3). The framework then foresees an integration function (level 4), before reaching the
393 Effectiveness Evaluation Committee (level 5) and the Conference of the Parties (level 6).

394 24. The flow of information and analysis is represented in Diagram 1 below, and explained in
395 more detail thereafter:



396
397
398
399 25. **Level 1 – Information:** As a first step, information resources available to support the
400 effectiveness evaluation will be identified and amassed. This will include information
401 from reports mandated by the Convention (e.g., implementation reports per Article 21;
402 compliance information and recommendations per Article 15; reports on effectiveness of
403 financial mechanisms per Article 13; emission inventories under Article 8 and release
404 inventories under Article 9; ASGM National Action Plans (NAP) progress reports under
405 Article 7), as well as the Secretariat’s Capacity Building and Technical Assistance report.

406 26. Such reporting may be incomplete, and thus these reports will be supplemented by other
407 available scientific, environmental, technical, financial and economic information per
408 Article 22 paragraph 3 and Article 19.

- 409 27. Clear criteria for this data collection should be established (e.g. including peer-reviewed
410 research articles and official publications such as national reports). The monitoring
411 arrangements are specified in Annex 3 and will determine which monitoring resources
412 will be included in the effectiveness evaluation.
- 413 28. **Level 2 – Compilation:** Relevant data for the effectiveness evaluation will be extracted
414 from the selected resources and compiled into a format that will enable their use in the
415 subsequent assessment and evaluation stages. Quality control of data should be conducted
416 at this stage. For monitoring data, this may include the compilation of monitoring data
417 into a global/central database with a consistent format, common quality control/quality
418 assurance procedures, assessment of confidence, and other relevant elements.
419
- 420 29. **Level 3 – Synthesis reports:** The amassed and compiled data will be used to create a set
421 of reports that synthesize the information. These reports are to inform the four policy
422 questions. Responding to the request from the Conference of the Parties that the
423 evaluation is to provide for an integrative picture of the general effectiveness of the
424 Convention, the ad hoc technical expert group lays out that the following synthesis
425 reports will be required (see Annex 4 for descriptions of the synthesis reports):
- 426 1. The state of global mercury levels in the environment, biota and humans, as well
427 as trends over time – **Global Monitoring Report**;
 - 428 2. **Emissions and Releases Report** – modelled after the *Global Mercury*
429 *Assessment* (2018);
 - 430 3. **Trade, Supply and Demand Report** – modelled after *Global Mercury: Supply,*
431 *Trade and Demand* (2017);
 - 432 4. **Waste Management Report** – building on *Global Mercury Waste Assessment*
433 (2017).
- 434 30. The reports will inform on the process, outcome and monitoring indicators, to facilitate
435 the Effectiveness Evaluation Committee’s consideration of the four policy questions.
436 These reports will include scientific and technical background, as well as accessible
437 visual presentations.
- 438 31. **Level 4 – Integrated Assessment Report:** The synthesis reports (and, where needed, the
439 underlying and/or additional data) on Convention actions, emissions and releases, and
440 monitoring data, etc. will be used for the fifth, the **Integrated Assessment Report**. This
441 integrated assessment report will distinguish between the process, outcome and
442 monitoring indicators to facilitate the Effectiveness Evaluation Committee’s efforts to
443 address the four policy questions. The report will take into account other information
444 (information such as socio-economic information, technology innovation, climate data,
445 key global policies, etc.) as necessary. (See annex 4 for a further description of the
446 Integrated Assessment Report).
- 447 32. The analyses will likely also include modelling to estimate how changes in emissions and
448 releases due to Convention measures have contributed to changes in mercury levels in the
449 environment, humans and biota.

Deleted: 8

- 451 33. The integrated assessment will also seek to evaluate long-term trends to interpret the
452 relevance of social, technical and economic data in the context of effectiveness the
453 Convention vis-à-vis its objective.
- 454 34. While the Integrated Assessment Report is to be an evidence-based science and technical
455 report, it is also to be accessible to non-technical readers and include visual
456 representations.
- 457 35. **Level 5 – Effectiveness Evaluation:** The Integrated Assessment Report supplemented,
458 as necessary, by the above-mentioned synthesis reports, will be submitted to the
459 Effectiveness Evaluation Committee. The Committee will use this information to
460 consider the four policy questions to derive conclusions about the effectiveness of the
461 Convention. The Committee may include in its report suggestions for improving the
462 effectiveness evaluation framework. The Committee may also highlight areas that the
463 Conference may wish to consider for the effectiveness of the Convention.
- 464 36. **Level 6 – Conference of the Parties:** The Conference of the Parties receives and reviews
465 the report of the Effectiveness Evaluation Committee and considers the conclusions and
466 recommendations of the Committee. The Conference makes its determinations regarding
467 actions or mechanisms to improve the effectiveness of the Convention.

468
469 **2. Development of indicators**

- 470 37. The Minamata Convention includes a number of measures that seek to control, reduce or
471 eliminate the major sources and uses of mercury, as well as a set of further stipulations
472 that oblige Parties to work together to support each other in the overall endeavour to
473 protecting people and the environment from the adverse effects of mercury.
- 474 38. To provide an integrative picture of the general effectiveness of the measures and
475 provisions of the Convention, the ad hoc technical expert group used an integrative
476 approach to identify indicators.
- 477 39. A set of indicators on process, outcome and monitoring to inform the policy questions are
478 proposed. The proposed indicators draw on previous work on elements for the
479 effectiveness evaluation framework, and the review of monitoring capacities and abilities.
- 480 40. Numerous indicators, developed following an article-by-article review, are clustering to
481 enable synthesised analysis in the proposed reports. The following articles and indicators
482 are clustered for evaluative purposes:
- 483 (a) **Supply cluster:** supply, storage and waste (Art 3, 10, 11);
484 (b) **Demand cluster:** products, processes and ASGM (Art 4,5,7);
485 (c) **Pressures cluster:** emissions, releases and contaminated sites (Art 8, 9, and 12),
486 (d) **Support cluster:** financial and technical assistance (Art 13 and 14), and
487 (e) **Information and research cluster:** information exchange, public information,
488 research (Art 17, 18 and 19).
- 489 41. Furthermore, recognising the central nature of some articles, such as Article 1 (setting out
490 the objective of the Convention) or the crosscutting nature, such as Article 16 (on the

491 health aspects) key articles were not clustered but kept separate for the purposes of
492 identifying indicators.

493 42. The rationale underlying the proposed indicators is as follows: (a) Process indicators are
494 required to answer the first policy question (Have the Parties taken actions to implement
495 the Minamata Convention?). (b) Outcome indicators are required to address the second
496 policy question (Have the actions resulted in changes in emission and releases of mercury
497 to the environment?). For each cluster of articles, the ad hoc expert group followed the
498 formulation of identification of how many parties are taking action on a key policy
499 measure, and what is the outcome of those actions. (c) Monitoring indicators are needed
500 to provide validated, scientific information to inform and support policy and decision-
501 making.

502 43. The indicators (a) and (b) above were largely developed keeping in mind data and reports
503 required by the Convention's reporting requirements or related bodies (including, for
504 example, reports of the Global Environment Facility). These reports will be
505 supplemented by other available and compiled data in the synthesis reports, and in the
506 Integrated Assessment Report. By using the data available, the indicators are therefore
507 cost effective. Further, the data will be produced on a recurrent basis for the life of the
508 Convention, and thus are sustainable.

509 44. The indicators are formulated in a way that can be practical and feasible. The indicators
510 are designed to be easily counted and calculated, and to be easily understood (they do not
511 represent complex functions). If Article 21 reporting data is submitted electronically to
512 the Secretariat, their calculation should be especially straight forward.

513 45. Baselines are considered fundamental to undertake an effectiveness evaluation, so that
514 indicators can be evaluated over time. There is no formal process under the Convention
515 to establish baselines. There are two approaches to establish baselines. One is a "before-
516 after" baseline, another is "with-without" baseline. The former is suitable for the
517 indicators that are relatively stable, so that a time value from before the Convention can
518 be used throughout the evaluation process. The latter type is suitable for indicators that
519 fluctuate over time by some factors other than the interventions made due to the
520 provisions of the Convention. Socio-economic and demographic aspects can play a role,
521 as can climate change, ongoing initiatives, as well as shifts in life style. These will impact
522 baseline value in the medium and longer term.

523 46. Table 2 below presented the proposed indicators, that are to be read in compliment to the
524 specific monitoring indicators identified in paragraph 52:

525

Commented [m2]: Insertion suggested: Indicator (c) above is not relevant to this paragraph.

Table 2: Proposed indicators to evaluate the effectiveness of the Minamata Convention			
A: Minamata Convention Article 1: (Objective) Protecting human health and the environment **		Source of information on indicator	Baseline for the indicator
A1. Cross-cutting monitoring indicator	Levels of mercury in the environment and in humans due to anthropogenic emissions and releases	- Integrated modelling	Baseline amount in the first evaluation (if models are available)
Notes	<ul style="list-style-type: none"> ▪ Attribution to be estimated using modelling to be developed ▪ In case of non-availability of such information from models, levels of mercury and trend in mercury (changes over time) will be used. ▪ The indicator for Article 1 is to be read with the relevant monitoring indicator indicated in Table 4, paragraph 52. 		

B: Supply Cluster of Articles: Mercury supply sources and trade (Article 3), Environmentally sound interim storage of mercury other than waste mercury (Article 10), Mercury waste (Article 11)		Source of information on indicator	Baseline for the indicator
B1. Overall process indicator for Articles 3, 10 and 11	Share of Parties that have implemented key provisions under this cluster (encompassing all process indicators below, i.e. B5, B6, B7, B9 and B12)	- Article 21 reporting	Baseline amount in the first evaluation
B2. Additional Cross-cutting outcome indicator for Articles 3, 10 and 11	Estimated global supply of mercury, in tonnes per year	- Synthesised information from individual indicators for Art 3, 10 and 11	Baseline amount in the first evaluation
Article 3			
B3. Outcome indicator for Article 3	Total amount of Hg mined from primary mercury mines	- Global Mercury Trade, Supply, Demand (2017) - ASGM NAP reports	Baseline amount in the first evaluation
B4. Outcome indicator for Article 3	Amount of Hg traded - broken down for specific purposes	- Article 3 forms	Baseline amount in the first evaluation
B5. Process indicator for Article 3	Number of parties that have developed an inventory of stocks and sources of supply	- Article 21 reporting	Baseline number in the first evaluation
B6. Process indicator for Article 3	Share of parties that have excess Hg from Chlor Alkali that have taken measures that such mercury is subject to final disposal	- World Chlorine Council Reports	Baseline % in the first evaluation
B7. Process indicator for Article 3	Number of parties trading in mercury	- Article 3 forms	Baseline amount in the first evaluation
Article 10			

B9. Process indicator for Article 10	Number of parties that have taken measures to ensure sound interim storage	- Article 21 reporting	Baseline amount in the first evaluation
B10. Outcome indicator for Article 10	Amount of Hg stored in an environmentally sound way as identified in the inventory of stocks	- Article 21 reporting	Baseline amount in the first evaluation
Article 11			
B11. Outcome indicator for Article 11	Amount of mercury/mercury compound waste subjected to final disposal	- Article 21 reporting	Baseline amount in the first evaluation
B12. Process indicator for Article 11	Number of parties that have measures in place to manage mercury waste in an environmentally sound manner	- Article 21 reporting	Baseline amount in the first evaluation
Notes	<ul style="list-style-type: none"> ▪ Data from non-Parties is important too. 		

C: Demand Cluster of Articles: Mercury-added products (Article 4), Manufacturing processes in which mercury or mercury compounds are used (Article 5), and Artisanal and small-scale gold mining (Article 7)		Source of information on indicator	Baseline for the indicator
C1. Cross-cutting process indicator for Articles 4, 5 and 7	Share of Parties that have implemented key provisions under this cluster	- Synthesised information from individual indicators for Art 4, 5 and 7	Baseline % in the first evaluation
C2. Cross-cutting outcome indicator for Articles 4, 5 and 7	Global use of Hg product or process in tonnes per application	- Information from industry stakeholders	Baseline amount in the first evaluation
Article 4			
C3. Process indicator for Article 4	Number of parties having appropriate measures to not allow the manufacture, export or import of mercury-added products listed in Part I of Annex A	- Article 21 reporting	Baseline number in the first evaluation
C4. Process indicator for Article 4	Number of exemptions per product category which are still valid	- Registry of exemptions	Baseline number in the first evaluation
C5. Process indicator for Article 4	Number of parties that have taken two or more measures for the mercury-added products listed in Part II of Annex A	- Article 21 reporting	Baseline number in the first evaluation
C6. Additional outcome indicator for Article 4	Volume tonnes of mercury added products (a) imported and (b) exported, in units per year for each product category in Annex A Part I	- Trade and customs data	Baseline amount in the first evaluation
Article 5			
C7. Process indicator for Article 5	Number of parties with exemptions for Annex B Part 1 processes, which are still valid	- Registry of exemptions	Baseline number in the first evaluation

C8. Process indicator for Article 5	Number of parties having measures in place to not allow the use of mercury or mercury compounds in manufacturing processes listed in Part I of Annex B	- Article 21 reporting	Baseline number in the first evaluation
C9. Process indicator for Article 5	Share of the parties that have processes subject to Article 5 para 3, that have taken all the measures for the respective processes listed in Annex B, Part II	- Article 21 reporting	Baseline % in the first evaluation
Article 7			
C11. Outcome indicator for Article 7	Total amount of Hg used in ASGM globally, in tonnes per year	- Article 21 reporting - NAPs and its review - Notifications	Baseline amount in the first evaluation
C12. Process indicator for Article 7	Share of parties declaring more than insignificant ASGM that have submitted NAP	- Notifications	Baseline % in the first evaluation
C13. Process indicator for Article 7	Share of parties that have submitted a NAP and have reviewed it	- Article 7 review	Baseline % in the first evaluation
Notes	▪ Some data on products may not be obtainable from public sources.		

D: Pressure Cluster of Articles: Emissions (Article 8), Releases (Article 9) and Contaminated Sites (Article 12)		Source of information on indicator	Baseline for the indicator
D1. Overall process indicator for Articles 8, 9 and 12	Share of Parties that have implemented key provisions under this cluster	- Article 21 reporting	Baseline % in the first evaluation
D2. Cross-cutting outcome indicator for Articles 8, 9 and 12	Total amount of Hg emitted and released	- Global Mercury Assessment - Inventories - MIAs	Baseline amount in the first evaluation
Article 8 **			
D3. Outcome indicator for Article 8	Total amount of Hg emitted from each of point source categories in Annex D (Article 21 report, inventories)	- Article 21 reporting	Baseline number in the first evaluation
D4. Process indicator for Article 8	Number of parties that have enacted appropriate laws and regulations to require BAT/BEP for new sources	- Article 21 reporting	Baseline number in the first evaluation
D5. Process indicator for Article 8	Number of parties that have put in place control measures for existing sources (per each of the measures set out in Article 8, para 5)	- Article 21 reporting	Baseline number in the first evaluation
Article 9 **			

D6. Outcome indicator for Article 9	Total amount of Hg releases in the inventory from relevant sources (Article 21 report, inventories)	- Article 21 reporting	Baseline number in the first evaluation
D7. Process indicator for Article 9	Number of parties that have identified relevant sources	- Article 21 reporting	Baseline number in the first evaluation
D8. Process indicator for Article 9	Number of parties that have established inventory of releases from relevant sources	- Article 21 reporting	Baseline number in the first evaluation
Article 12			
D9. Process indicator for Article 12	Number of parties that have developed strategies for identifying and assessing sites contaminated by mercury or mercury compounds	- Article 21 reporting	Baseline number in the first evaluation
D10. Process indicator for Article 12	Number of parties that have developed the inventory of contaminated sites	- Article 21 reporting	Baseline number in the first evaluation
Notes	<ul style="list-style-type: none"> ▪ The indicators for Article 8 and 9, are to be read with the relevant monitoring indicators indicated in Table 4, paragraph 52. 		

E: Support Cluster of Articles: Financial resources and mechanism (Article 13), and Capacity-building, technical assistance and technology transfer (Article 14)		Source of information on indicator	Baseline for the indicator
Article 13			
E1. Process indicator for Article 13	Number of Parties: <ul style="list-style-type: none"> ○ that have contributed to the financial mechanism referred to in paragraph 5 of Article 13 ○ that have received GEF resources ○ that have received SIP resources ○ that have mobilised national resources for implementing the Convention 	- Article 21 reporting	Baseline number in the first evaluation
E2. Process indicator for Article 13	Amount of resources provided by: <ul style="list-style-type: none"> ○ Global Environment Facility ○ Specific International Programme ○ Bilateral support 	- Article 21 reporting	Baseline number in the first evaluation
E3. Additional Process indicator for Article 13	Number of recommendations from the financial review reflected in the GEF/SIP policy documents	- Information from policy documents	Baseline: zero
Article 14			
E4. Process indicator for Article 14	Number of Parties: <ul style="list-style-type: none"> ○ that have cooperated for providing capacity building and technical assistance to another party 	- Article 21 reporting	Baseline number in the first evaluation

	<ul style="list-style-type: none"> ○ that have requested technical assistance ○ that have received capacity building or technical assistance ○ that have promoted or facilitated technology transfer 		
Notes	<ul style="list-style-type: none"> ▪ The cycle of review of the Financial Mechanism may well not align with the effectiveness evaluation cycle. 		

530

F: Minamata Convention Article 15: Implementation and Compliance Committee		Source of information on indicator	Baseline for the indicator
F1. Process indicator	Proportion of issues that the Committee was able to resolve, including indications of systemic issues, if any	- ICC report, as referred to in Art 21	Baseline number in the first evaluation
Notes	<ul style="list-style-type: none"> ▪ The expert group could not complete its deliberations in the indicator, as the ICC had not yet finalised its terms of reference. Their terms of reference are to be considered by COP3. 		

531

G: Minamata Convention Article 16: Health aspects **		Source of information on indicator	Baseline for the indicator
G1. Monitoring indicator	Mercury levels in selected human populations (as defined by the monitoring arrangements)	- Existing monitoring data and activities	Baseline number in the first evaluation
G2. Process indicator	Number of parties that have taken measures, such as fish advisories, to provide information to the public on exposure to mercury in accordance with paragraph 1 of article 16	- Article 21 reporting	Baseline number in the first evaluation
G3. Process indicator	Number of parties that have taken measures to protect human health in accordance with article 16	- Article 21 reporting - Submissions to the secretariat	Baseline number in the first evaluation
Notes	<ul style="list-style-type: none"> ▪ The indicator for Article 16 is to be read with the relevant monitoring indicators indicated in Table 4, paragraph 52. ▪ Mercury levels in biota also to be considered. 		

532

H: Information and Research Cluster of Articles: Information exchange (Article 17), Public information, awareness and education (Article 18), Research, development and monitoring (Article 19)		Source of information on indicator	Baseline for the indicator
Article 17			
H1. Process indicator for Article 17	Number of parties with designated national focal points	- Article 21 reporting	Baseline number in the first evaluation
H2. Process indicator for Article 17	Number of parties that are participated in information exchange mechanisms related to mercury	- Submissions to the secretariat	Baseline number in the first evaluation
Article 18			

Deleted: have established

Commented [m3]: Revision suggested: Information exchange is inter-agency mechanism, so each party does not have to 'establish' a mechanism by itself.

H3. Process indicator for Article 18	Number of parties that have taken measures to implement article 18	- Article 21 reporting	Baseline number in the first evaluation
H4. Process indicator for Article 18	Average number of measures under paragraph 1 of Article 18 that are being implemented by parties	- Derived from Article 21 reporting	Baseline number in the first evaluation
H5. Process indicator for Article 18	Number of parties that have public information on mercury levels in air, humans and biota	- Article 21 reporting	Baseline number in the first evaluation
H6. Process indicator for Article 18	Number of parties undertaking risk communication relating to mercury intake through food and water consumption	- Article 21 reporting	Baseline number in the first evaluation
Article 19			
H7. Process indicator for Article 19	Number of parties that have undertaken research, development and monitoring in accordance with paragraph 1 of article 19	- Article 21 reporting	Baseline number in the first evaluation
H8. Process indicator for Article 19	Number of parties contributing data and knowledge to integrated assessments	- Existing monitoring networks, databases, scientific data and literature	Baseline number in the first evaluation
H9. Additional process indicator for Article 19	Number of regions contributing to a regional dataset	- Existing monitoring networks, databases, scientific data and literature	Baseline number in the first evaluation
Notes	<ul style="list-style-type: none"> Submissions to the Secretariat that supplement article 21 reporting. 		

Commented [m4]: Revision suggested: Mercury is not consumed but enters people's body through food and water.

534

I: Minamata Convention Article 20: Implementation Plans		Source of information on indicator	Baseline for the indicator
I1. Process indicator	Number of parties submitting implementation plans	- Secretariat report to the COP submissions	Baseline: zero
Notes	<ul style="list-style-type: none"> Parties do not have the obligation to prepare implementation plans. Some Parties found it useful to prepare such a plan nevertheless, and submit it to the Secretariat. 		

535

J: Minamata Convention Article 21: Reporting		Source of information on indicator	Baseline for the indicator
J1. Process indicator	Proportion of parties reporting on time	- Article 21 reporting	Baseline: % of the first submission on time
J2. Process indicator	Proportion of parties indicating that information is not available for specific questions	- Article 21 reporting	Baseline: % not available in the first reports

Notes	<ul style="list-style-type: none"> ▪ Parties are to report every two years.
--------------	--

536

K: Minamata Convention Article 22: Effectiveness evaluation		Source of information on indicator	Baseline for the indicator
K1. Process indicator	Evidence of implementation of recommendations from effectiveness evaluation through decisions and actions of the Conference of the Parties	- COP report	Baseline: zero
Notes	<ul style="list-style-type: none"> ▪ This article will not be evaluated in the first evaluation. 		

537

538

539

540 **3. Data sources**

541 47. The availability of information sources from which to derive these indicators, as well as
 542 supplementary relevant and comparable scientific, environmental, technical, financial and
 543 economic information on which to base the effectiveness evaluation, is driven by a
 544 number of factors.

545
 546 48. First, different articles of the Convention have different time lines for implementation.
 547 Some of these deadlines fall within the first cycle of effectiveness evaluation (2017-
 548 2023), but some do not. Moreover, even if a measure is implemented within the deadline,
 549 evidence of its impact and therefore effectiveness may not be available for some time or
 550 not directly measurable. This presents some challenges on how to attribute effect. Table 3
 551 below gives a short overview of phase-out dates and time-bound articles of the Minamata
 552 Convention and their time lines.
 553

Table 3: Minamata Convention Timelines		
Date	Article	Description
2018	Art 5, para 2, Annex B	Acetaldehyde production in which mercury or mercury compounds are used as a catalyst to be phased out
2020	Art 4, para 1, Annex A	Manufacture, import or export of various mercury-added products shall not be allowed (including of batteries, switches and relays, compact and linear fluorescent lamps, high pressure mercury vapour lamps, cold cathode fluorescent lamps and external electrode fluorescent lamps for electronic displays, cosmetics, pesticides and topical antiseptics, as well as barometers, hygrometers, manometers, thermometers and sphygmomanometers)
	Art 5, para 3, Annex B	In vinyl chloride monomer production, reduce the use of mercury in terms of per unit production by 50 per cent by the 2020 against 2010 use. For sodium or potassium methylate or ethylate reduce emissions and releases in terms of per unit production by 50 per cent by 2020 compared to 2010
2025	Art 5, para 2, Annex B	Chlor-alkali production to be phased out
2027	Art 5, para 3, Annex B	For sodium or potassium methylate or ethylate, reduce the use of mercury aiming at phase-out of its use as fast as possible and within 10 years of entry into force of the Convention
2035	Art 3, para 4	Primary mining of mercury that was conducted within a Party's territory at the date of entry, is to cease 15 years after that date
2020	Art 5	Submit to the secretariat information on the number and types of facilities covered under Annex B, and the amount of mercury or mercury compounds used
	Art 7	Submit NAP to secretariat if developed
	Art 9	Identify relevant point source categories
2021	Art 8 and 9	Submit National Implementation Plan if one has been developed

2022	Art 8	Require BAT/BEP for new facilities
	Art 8	Develop and maintain an inventory of emissions sources
	Art 9	Develop and maintain an inventory of release sources
2017	Art 8	Require measures for control on existing facilities

554

555 49. Second, various important identified data sets and information sources that have been
556 produced in the past are not required under the Convention, and thus the production of
557 similar reports in the future is not assured or governed by Convention requirements.
558 These include Minamata Initial Assessment, as well as the Global Mercury Assessment
559 (produced 2002, 2008, 2013 and 2018), and the reports on Global Mercury: Supply,
560 Demand and Trade (2006 and 2017).

561 50. Third, some information sources differ in frequency. The ASGM National Action Plans
562 due 3 years after the entry into force of the Convention for that party (or three years after
563 its declaration to the Secretariat that it has more than insignificant ASGM) and must be
564 reviewed every three years thereafter. The regular reports under Article 21 are to be
565 completed every two years for specified questions, with the first short reports due at the
566 end of 2019, and the first full reports that respond to all reporting requirements due at the
567 end of 2021.

568 51. Last, is the consideration of the availability of relevant and comparable scientific and
569 environmental monitoring data. On the one hand, mercury is one of the longest studied
570 chemicals. On the other hand, in considering the available monitoring information and
571 available data on mercury and mercury compounds, the ad hoc technical expert group
572 noted that not all data and matrices are suitable for direct comparison or analysis at the
573 global level, and modelling will be critical to shape our fuller understanding of the
574 presence of mercury and its trends in our environment, as well as to attribute changes in
575 mercury levels to Convention measures.

576
577 52. The current mercury monitoring arrangements and modelling frameworks are outlined in
578 more detail in Annex I on Technical Information on Monitoring and Annex 3 on the
579 Global Monitoring Arrangements. Table 4 below given an indicative overview of
580 selected monitoring indicators and how they can contribute to the overall effectiveness
581 evaluation. They are to be read in complement with the indicators presented in Table 2 in
582 paragraph 46:
583
584

Table 4: Monitoring indicators by media		
Overall media	Indicator	Source of info
1. Air	M1. Total gaseous mercury/gaseous elemental mercury in the ambient air	Existing/expanded monitoring activities and networks

	M2. Mercury level in precipitation	Existing/expanded monitoring activities and networks
2. Human	M3. Mercury level in hair as primary matrix	Epidemiological studies by Parties
	M4. Mercury level in blood as alternative	International and national biomonitoring programme Longitudinal birth cohort and cross-sectional studies
3. Biota	M6. Mercury levels in biota	Continental network
	M7. Mercury levels in biota	Oceanic framework
4. Water	M8. Mercury levels in sea water covering horizontal and vertical distribution	Existing/expanded monitoring activities and networks

Deleted: Water as a separate media is included to inform modelling (attribution)....

585
586

4. Use of modelling in the effectiveness evaluation

587 53. Models provide for the integration of mechanisms and observations and use that to assess
588 projections for future source apportionment. It can be said, that models therefore
589 formalise the scientific understanding of mechanisms affecting mercury behaviour. One
590 critical source of models is to provide a tool for linking and spatially/temporally
591 extrapolating monitoring data collected globally as part of ongoing research programmes,
592 policy activities and data provided by civil society, in order to provide a comprehensive
593 picture of the state of mercury pollution globally. Moreover, integrating modelling
594 frameworks provide a tool to work across media, i.e. for linking releases of mercury to
595 the atmosphere, land and water to methylmercury in fish and wildlife, as well as exposure
596 of human populations.

597 54. Another critical use of models in effectiveness evaluation is to attribute changes to levels
598 in mercury to Convention measures.

599 55. Models for different media (air, human, biota, water, and soil) vary in their ability and
600 state of development. For example, for air and atmosphere, many monitoring groups have
601 developed global modelling tools that can be used to assess the impact of changes in
602 anthropogenic mercury emissions and releases on global atmosphere concentrations, and
603 mercury inputs to terrestrial ecosystems and the ocean. Atmospheric models have been
604 extensively evaluated and can be applied to assess spatial gradients in atmospheric
605 mercury concentrations and deposition, as well as temporal changes. By contrast, models
606 for other media such as land, are still mainly used in research applications.

607
608 56. To bridge linkages across different media, integrated model frameworks seek to link
609 various models used for different media. In this way integrated modelling frameworks

612 provide a tool for linking emissions of mercury to the atmosphere and releases to land
613 and water, to methylmercury in fish and wildlife, and to exposure of some fish-
614 consuming human populations. It is to be noted that a difficult link in integrated
615 modelling frameworks is to human exposure and health outcomes due to the diversity of
616 dietary preferences, food consumption patterns and individual variability in
617 toxicokinetics affecting methylmercury uptake and elimination.

618 57. In addition to models that describe behaviour of mercury in the environment and
619 receptors, a variety of models and quantitative techniques can simulate socio-economic
620 systems to forecast where mercury is present in society and where it might eventually
621 enter the environment. In this way models can be used to develop scenarios that represent
622 baseline and different policy alternatives. Inputs to these models include commercial data
623 (e.g. amount of mercury in products), technological performance, economic information,
624 energy data, demographic information, policy specifications, and institutional analysis.
625 Outputs can include emissions and releases of mercury, and socio-economic parameters.
626 Other types of models that are relevant to understanding socio-economic systems of
627 relevance to mercury include life-cycle analysis, materials flow analysis, input-output,
628 and economic models.

629 58. Developing and evaluating integrated models draws on expertise that bridges natural
630 science, social science, and engineering. The components for an integrated modelling
631 framework are currently undergoing rapid development in the scientific and academic
632 community and should be available for our greater understanding of mercury cycling and
633 its impacts in the near future. It is expected such models will available by 2023 for
634 policy-relevant analyses.
635

636 5. Scientific and technical functions

637
638 59. The framework foresees two scientific and technical functions to be performed for the
639 effectiveness evaluation, namely a synthesis function, and an integration function. These
640 function at different levels of the framework.

641 a. **Synthesis function:** The first function is to synthesise mercury information
642 collected and compiled by the level 1 to 3 activities. Two streams of information
643 are referred to in this regard: (i) information provided by Parties based on Article
644 21 reporting, and (ii) information and knowledge that is scientific, peer-reviewed
645 and publicly available. The information is used to respond to the first three policy
646 questions, and the indicators identified, to prepare the four synthesis reports. This
647 function foresees a role for the secretariat, for scientists and experts, and for
648 organisations. This function foresees the production of four synthesis reports, one
649 of which is the Global Monitoring Report, for which a specific expert group is
650 foreseen.

651 b. **Integration function:** The second function, which occurs at level 4, is to interpret
652 the information and knowledge collected, compiled and synthesised by the level 1
653 to 3 activities to interpret the linkages between policy actions, emission reductions
654 and resulting mercury levels, using available data sources, modelling techniques
655 and analytical tools drawn from natural and social sciences. Further this function

656 included also the collection of additional non-mercury information for further
657 analysis that will include, but is not limited to, data on socio-economic and
658 demographic information such as global population trends, trade and industry
659 trends, mitigation and adaptation policies, or on technological innovation
660 information such as alternative products, communication and transport
661 technologies. The intent of this function is to provide an integrative picture
662 through the contextualisation of information for that evaluation cycle of the
663 Convention. The function foresees the production of the Integrated Assessment
664 Report, and the establishment of a specific integrated assessment group of
665 expertise.


666 60. The framework also differentiates between input from the following two:

667 a. **Monitoring:** Compiling, assessing and summarizing available monitoring data,
668 per the monitoring arrangements in Annex III, to describe the current state of
669 mercury concentrations, as well as trends in the environment, humans and biota,
670 and working with modelling experts as appropriate. The global monitoring report
671 developed at level 3 will be the input to the integrated assessment at level 4, and
672 also submitted to the Effectiveness Evaluation Committee.

673 b. **Modelling:** Analysing the contribution of emissions and releases covered by the
674 Convention to overall mercury concentrations in the environment, and where
675 possible, in humans and biota. Modelling conducted during level 4 will estimate
676 future mercury concentrations that reflect the overall impacts of mercury
677 emissions and releases, from legacy emissions and releases to those predicted in
678 the future under various scenarios, based on the reports made available in the
679 effectiveness evaluation process, as well as available relevant socio-economic
680 information.

681 **6. Institutional Arrangements for the Effectiveness Evaluation**

683 61. To implement the effectiveness evaluation process described thus far, the framework
684 identifies different entities that may deliver the tasks to compile, summarize and integrate
685 data and knowledge, and to perform relevant scientific and technical analyses. Diagram 2
686 below displays the activities, outputs and flow of information and knowledge among
687 entities potentially responsible for these tasks.



689 **Minamata Convention Secretariat**

- 690 62. The Secretariat will play a role in collecting, compiling, summarizing and synthesizing
691 available data. The Secretariat already has a role, prescribed by the Convention, to act as
692 the mechanism through which Parties submit reports under Article 21, which in turn will
693 contain references to progress reports on the NAPs, under Article 7, to inventories under
694 Articles 8 and 9; and voluntary NIPs under Article 20. The Secretariat may, as
695 appropriate, be assisted by groups of experts or hired experts, conduct literature reviews,
696 produce datasets for further analysis or organize synthesis and peer review.
- 697 63. These datasets will be processed at level 3 for calculating/ tabulating process and
698 outcome indicators. The Secretariat will also become responsible for facilitating synthesis
699 reports that combine these indicators with other relevant information, including
700 commissioning external expertise where necessary, as UNEP has done in previous efforts
701 – for example, under the Global Mercury Assessment (2018), the report on Global
702 Mercury Supply, Trade and Demand (2017) and the Global Mercury Waste Assessment
703 (2017).
- 704 64. The Secretariat will also compile summaries and synthesis reports resulting from other
705 processes mandated by the Convention, such as reports from the Implementation and
706 Compliance Committee under Article 15, reports from bodies implementing the financial
707 mechanism, the report on the effectiveness of the finance mechanism, required under
708 Article 13, paragraph 11 (which will draw inter alia on reports such as GEF report and
709 the SIP report) and the Secretariat’s report on Capacity Building and Technical
710 Assistance. All synthesis reports and summary documents will be eventually submitted to
711 the Effectiveness Evaluation Committee as supplementary information for their
712 consideration at level 5. These reports (and underlying data where needed) will be
713 transmitted for integrated assessment at level 4.
- 714 **Delivery of the scientific and technical functions**
- 715 65. The framework puts forward that the scientific and technical functions can be delivered
716 as follows:
- 717 a. **Scientific and Technical Expertise:** A scientific and technical grouping
718 comprising of individuals with extended expertise on monitoring, scientific and
719 technical assessment, and natural and social sciences and research relevant to
720 mercury, is to deliver the activities of level 1 to 3, to produce the four synthesis
721 reports. For this purpose, there are roles for the secretariat, for scientists and
722 experts, and for organisations. This group will include a specific group of
723 monitoring and modelling experts to coordinate monitoring and modelling
724 activities that produces the Global Monitoring Report (a synthesis report).
- 725 b. **Integration Assessment Group:** A small separate group is required, at level 4, to
726 produce the Integrated Assessment Report for the Effectiveness Evaluation
727 Committee. Specific chapter and section authors led by a chief author will be
728 identified to comprise this group. The group will necessarily be multi-disciplinary
729 in nature, and authors will be identified according to their most suitable expertise.
730 For attribution functions, the group will include modellers. Additionally, this
731 group will also be supported by communication expertise to ensure the results of

Deleted: 8

733 this integrated assessment are summarised and presented in visual forms (e.g. a
734 dashboard type score table summarising progress).

735 **Effectiveness Evaluation Committee**

736 66. The Effectiveness Evaluation Committee at level 5 will use the Integrated Assessment
737 Report supplemented by the four synthesis reports to consider the policy questions posed
738 in the framework, and from that derive conclusions about the effectiveness of the
739 Convention. The Effectiveness Evaluation Committee will formulate recommendations
740 aiming at improving the effectiveness of the Convention. The Committee may include in
741 its report suggestions for improving the effectiveness evaluation framework. Terms of
742 reference for the Committee are found in Annex II.

743 **Conference of the Parties**

744 67. The ultimate responsibility for evaluating the effectiveness lies with the Conference of
745 the Parties at Level 6. In this framework, the intention is for the Conference to consider
746 the recommendations of the Committee, and then make determinations about any needed
747 changes to or strengthening of Convention measures. The Conference may also choose to
748 mandate changes in the procedures for future effectiveness evaluation cycles.

749 **Additional proposal to deliver scientific and technical functions**

750 68. The implementing structure for the scientific and technical functions can also be
751 delivered by an external entity following a bidding process. In this case, the Secretariat
752 could be asked to call for proposals that include, but is not limited to, the approach to
753 complete necessary tasks, structure to implement these tasks, associated costs, etc.
754 Should an entity for delivery of this function be selected through a bidding process, full
755 information on the process will be reported to the Conference of the Parties.
756
757
758

Deleted: expert

Deleted: expert

761 **7. Schedule and timetable**


762 69. Paragraph 1 of Article 12, Effectiveness Evaluation, holds that the Conference of the
 763 Parties shall evaluate the effectiveness of the Convention no later than six years after the
 764 entry into force of the Convention, and periodically thereafter at intervals to be decided
 765 on.

766 70. As the Convention entered into force on 16 August 2017, the outcome of the first cycle of
 767 evaluation it to be submitted in 2023. The fifth meeting of the Conference of the Parties
 768 will convene in that year.

769 71. Taking into account the four-year cycle of reports under Article 21 (Reporting), the
 770 regular reports due to the Conference of the Parties on Article 13 (Financial resources and
 771 mechanism) and Article 14 (Capacity-building, technical assistance and technology
 772 transfer), as well as the monitoring data available, the date line for comparable
 773 information for this first cycle of evaluation is recommended to be set as 2021.

774 72. This allows 2022 and 2023 to be utilised by the Effectiveness Evaluation Committee for
 775 its review and analysis to be finalised to be presented to the fifth meeting of the
 776 Conference of the Parties.

777 73. The timeline for the first cycle of the effectiveness evaluation of the Convention is set out
 778 in Diagram 3 below:



First Cycle of Effectiveness Evaluation of the Minamata Convention from 2017 to 2023

781 74. As the Conference of the Parties from its fourth meeting onwards, is to convene every
 782 two years, the future effectiveness evaluations would follow this six-year cycle.
 783
 784
 785

786 **IV. Issues for further considerations**

- 787
- 788
- 789 75. The ad hoc technical expert group proposes a framework for effectiveness evaluation that
- 790 follows a flow of information from level 1 to level 6, and identifies different entities that
- 791 fulfil different functions and roles in the process. While some of these entities already
- 792 exist (i.e. those for administrative and programme support, compilation of data for
- 793 synthesis reports, etc), there are others who are to perform vital scientific and technical
- 794 functions to implement the framework, that are not in place yet.
- 795 76. The framework foresees two scientific and technical functions: (i) to produce four
- 796 synthesis reports (one of which is the Global Monitoring Report), and thereafter, (ii) to
- 797 produce the integrative picture (the Integrated Assessment Report). These reports are to
- 798 inform the deliberations of the Effectiveness Evaluation Committee, which in turn reports
- 799 the outcome of its evaluation to the Conference of the Parties.
- 800 77. To operationalise the all constituent elements of the framework, the Conference of the
- 801 Parties will need put a number of entities into place. Most entities conducting the
- 802 activities at the different levels are identifiable. They include, the Monitoring and
- 803 Modelling Group (which is to produce the Global Monitoring Report), the Integration
- 804 Assessment Group (which is to produce the Integrated Assessment report), and the
- 805 Effectiveness Evaluation Committee (which is to present its evaluation report to the
- 806 Conference of Parties). These can be put into place by the Conference of the Parties.
- 807 78. What is still to be clarified by the Conference of the Parties is which entities will produce
- 808 the following reports: (i) Emissions and Releases Report, (ii) Trade, Supply and Demand
- 809 Report, and (iii) Waste Management Report.

810

811

812 **Suggested action by the Conference of the Parties**

813 79. The Conference of the Parties may wish to consider the recommendations of the ad hoc
814 expert group on the proposed framework for the effectiveness evaluation, and may wish
815 to adopt a decision along the following lines:

The Conference of the Parties,

Welcoming the report on the proposed effectiveness evaluation framework and monitoring arrangements under the Minamata Convention on Mercury, and complementing information developed by the ad hoc technical expert group on the basis of mandates provided in decision MC-1/9 and decision MC-2/10;

Welcoming monitoring activities already in place and efforts of Parties and others to support the provision of monitoring data on mercury and their availability in the future;

Acknowledging the available modelling capacities for the use in the effectiveness evaluation;

1. *Adopts* the framework for the effectiveness evaluation proposed for the Minamata Convention, including its methodology, indicators, reports, schedule and timeline;
2. *Adopts* the terms of reference and mandate of the Effectiveness Evaluation Committee to perform the evaluation;
3. *Decides* to establish the committee at its fourth meeting;
4. *Establishes* a [scientific and technical group] [monitoring and modelling group] to perform the scientific and technical functions that enable the timely production of the required synthesis reports by its fourth meeting, to enable the integrative work to be done for the Effectiveness Evaluation Committee to convene ahead of its fifth meeting to complete the first evaluation of the Minamata Convention;
5. *Requests the Secretariat* to support the work of the effectiveness evaluation, and to continue to collect information relevant to the effectiveness evaluation including for the development of synthesis reports, working with relevant experts and organizations;
6. *Requests* the group on monitoring and modelling to work in line with its terms of reference including the finalization of monitoring guidance and develop a global monitoring report by COP4 to support the first effectiveness evaluation;
7. *Encourages* Parties to engage actively in the implementation of the effectiveness evaluation framework, in particular, to:
 - a) Continue to monitor mercury and to share the resulting monitoring data through the group established pursuant to paragraph 3;

- b) Collaborate to develop and improve modelling as needed, and carry on geographically representative monitoring of mercury in the environment, in humans, and in biota;
- c) Use tools supporting data harmonization identified such as standard operating procedures (SOPs), methodologies and techniques identified by the ad hoc expert group; and
- d) Support the further development and long-term implementation of the global monitoring arrangements, if in a position to do so.

816

817

818 **Annex 1: Technical information on monitoring**

819

820 **1. Introduction**

- 821 1. This annex summarizes the work done by the ad hoc group with regard to global
822 monitoring arrangements at its two meetings in March 2018 and April 2019, and through
823 electronic communication.
- 824 2. It starts in Section 2 with the identification of categories of the available comparable
825 monitoring data most effective in providing information on global trends, monitoring data
826 in air, water, biota, and humans that could be used to assess the impact on levels and
827 trends of mercury, and the potential and limitations of the data identified. Section 3
828 further assesses the extent to which the information reviewed meets the needs for
829 effectiveness evaluation, identifies major gaps, outlines options to enhance the
830 comparability and completeness of the information, and compares these options for their
831 cost-effectiveness, practicality, feasibility and sustainability, global coverage, and
832 regional capabilities to identify opportunities for future enhancements to monitoring.
833 Section 4 identifies available modelling capabilities to assess changes in global mercury
834 levels within and across different media. Section 5 examines options and identifies
835 sources of data that can be used for establishing a baseline for monitoring data. Further
836 discussion on the development of guidance for monitoring and proposed monitoring
837 arrangements is included in Annex 3 on terms of reference for global monitoring
838 arrangements.
- 839 3. A large amount of other relevant technical information on monitoring complementing the
840 proposal in this annex including an overview of available monitoring information, is
841 available in a reference document as UNEP/MC/COP.3/INF/xx.

842

843 **2. Identification of monitoring information/data**

844 **How monitoring activities may contribute to the development of the effectiveness**
845 **evaluation framework**

- 846 4. In considering monitoring information and data, the ad hoc group considered matrices
847 outlined in MC-2/9: air, biota, humans and water. The ad hoc group concluded that data
848 on levels of mercury and mercury compounds in air, biota and humans either are available
849 or would be able to be obtained, and would be comparable on a global basis. Some experts
850 were of the opinion that data on water are available on a global basis to some extent. The
851 availability and comparability of monitoring data for each matrix are discussed below.
- 852 5. Mercury levels in the atmosphere is directly linked to the emissions from the
853 anthropogenic sources identified by the Convention. The atmospheric monitoring
854 activities will contribute to the evaluation of the effectiveness of the Convention by
855 determining whether the levels of mercury are increasing or decreasing in the atmosphere
856 as per changes in the emissions of mercury and enable the modelling results to define
857 source-receptor relationships. Also, this data will contribute to the predictive capabilities
858 of regional and global models of mercury impacting the environment, which may also be
859 affected by other atmospheric chemistry issues.

860 6. Human biomonitoring has the following advantage in contributing to the effectiveness
861 evaluation of the Convention: provides information on exposure to mercury from all types
862 of sources; integrates the results of the different types of risk reduction measures, and
863 provides information on geographical distribution enabling identification of areas and
864 population groups requiring urgent support in terms of risk reduction measures.

865 7. Biota monitoring has an advantage in contributing to the effectiveness evaluation of the
866 Convention by tracking changes of environmental mercury levels at regional and global
867 levels to determine protection of human health and the environment.

868 **Ambient air**

869 8. Mercury levels in ambient air have been measured in some locations for a very long
870 period. These data have contributed to the discussion on the global nature of the mercury
871 issue. The current available data is collected by various national and global network
872 owners using different sampling methods. It was recognized that none of the currently
873 available data had global coverage, but that there are potential suitable methods to obtain
874 such global data (as identified in GMA 2018). Overview of existing networks is available
875 in the resource document (UNEP/MC/COP.3/INF/XX).

876 9. A number of suitable methods are available, and the available sampling techniques
877 considered suitable to obtain globally comparable data were identified and reviewed.
878 These include:

879 ▪ Total Gaseous Mercury (TGM) or Gaseous Elemental Mercury (GEM)
880 concentrations in air at background and impacted sites;

881 ▪ Wet deposition.

882 9. TGM/GEM can be measured adopting active continuous monitoring, manual active air
883 sampling and passive air sampling techniques. Active continuous techniques are in use at
884 several sites of existing regional and global monitoring networks and provide continuous
885 TGM/GEM concentrations, whereas manual active and passive sampling are used in
886 locations where no monitoring infrastructure is available and provide average TGM
887 concentrations as monthly (or at lower frequency) average.

888 10. The atmospheric deposition flux of mercury is considered the combination of wet and dry
889 deposition of mercury to the surface. Measurements of wet deposition are done through
890 the collection of rain samples and dry deposition either mathematically inferred or
891 measured through tree debris. Several existing long-term networks collect wet deposition
892 samples but, due to a lack of comparable standard procedures, dry deposition is not always
893 measured. The amount of total mercury measured in atmospheric deposition samples is
894 used as basis to calculate the total atmospheric deposition flux associated to a precipitation
895 (rain or snow) event.

896 11. Validated atmospheric mercury models are needed to assess source-receptor relationships
897 and evaluate the relative importance of each anthropogenic source and/or emission source-
898 region in the global mass balance of mercury with changing mercury emission regime,
899 meteorological conditions and climate forcing. Good global coverage of monitoring data
900 of mercury in ambient air and deposition samples are also of fundamental importance to
901 validate these atmospheric models. Further details are provided in
902 UNEP/MC/COP.3/INF/XX.

903 **Human exposure**

- 904 12. All people are exposed to some amount of mercury. For many communities worldwide,
905 dietary consumption of fish, shellfish, marine mammals, and other foods is arguably the
906 most important source of methylmercury exposure. Exposures to elemental and inorganic
907 mercury mainly occur in occupational settings (including artisanal and small-scale gold
908 mining) or via contact with products containing mercury. There remains high concern for
909 vulnerable groups including various indigenous populations with high dietary or
910 occupational exposure to mercury.
- 911 13. Human biomonitoring to assess general population exposure to mercury (i.e. background
912 level rather than “hot spots”) provides information on global trends. In the general
913 population, assessment of prenatal exposure is recommended because the fetus is the most
914 vulnerable to methyl mercury exposure.
- 915 14. There are two main biomarkers:
- 916 ▪ Total mercury in maternal scalp hair (3 cm hair strand from the scalp, to measure
917 exposure during the 3rd trimester).
 - 918 ▪ Total mercury in cord blood.
- 919 15. Scalp hair is a preferable biological matrix. It is easily available, a non-invasive method,
920 and there are no specific requirements for transportation and storage.
- 921 16. Cord blood can be alternative matrix to hair. Inclusion of cord blood in a survey provides
922 several additional advantages such as: demonstration of pre-natal exposure to mercury
923 (cord blood analysis characterizes both exposure of a mother and a child to mercury
924 during pregnancy); possibility to get more reliable results and exclude influence of
925 external factors (e.g. external contamination of hair by mercury, permanent hair treatment
926 decreasing mercury in hair); being an alternative biological matrix to hair in locations
927 where hair sampling is difficult due to cultural, ethical, religious specificities.
- 928 17. There are reliable, although variable, coefficients allowing comparability of results from
929 the mercury measurements in hair and blood/cord blood.
- 930 18. Assessment of total mercury is sufficient for characterizing exposure, unless external
931 exposure of scalp hair needs to be evaluated.
- 932 19. In addition to general population exposure, parties may conduct biomonitoring in other
933 vulnerable populations including the occupationally exposed and in hot spot areas. These
934 data may provide additional information of use for effectiveness evaluation, for example
935 when repeated over time in the same populations.
- 936 20. The Global Mercury Assessment 2018 has identified currently available data on mercury
937 exposure in regional and national human biomonitoring programmes, longitudinal birth
938 cohort studies and cross-sectional information in specific populations including high
939 exposure groups.
- 940 ▪ In regional and national human biomonitoring programmes, some information may be
941 comparable (depending on the ability to disaggregate data by sex and age within the
942 programme). Such studies are only available in a very small number of countries,

- 943 primarily in the northern hemisphere. Such studies are expensive and therefore not
 944 feasible for the sole purpose of monitoring global mercury exposure.
- 945 ▪ Comparable and high-quality data exists from a number of longitudinal birth cohort
 946 studies, including in groups consuming large amounts of seafood, freshwater fish
 947 and/or marine mammals. These are available only in a small number of locations,
 948 and are not globally representative.
 - 949 ▪ The GEF-funded project “Development of a Plan for Global Monitoring of Human
 950 Exposure to and Environmental Concentrations of Mercury” has generated
 951 comparable data in a small number of additional countries, using the WHO protocol.⁵
- 952 21. Total mercury in urine is relevant for populations with high exposure to elemental and
 953 inorganic mercury, and is not appropriate for assessment of methylmercury exposure. It
 954 may be useful for monitoring the impact of control actions taken by parties on mercury
 955 exposure in mining communities.
- 956 22. Human biomonitoring has a number of advantages for informing an assessment of the
 957 effectiveness evaluation of the Minamata Convention, including:
- 958 ▪ Directly addressing the fundamental question as to whether enough is being done to
 959 protect human health (Article 1 of the Convention);
 - 960 ▪ Integrating information on exposure to mercury from different sources;
 - 961 ▪ Integrating the effects of the range of risk reduction measures taken.
- 962 23. In using human biomonitoring data, it should be noted that human mercury level is
 963 affected by many confounding factors such as fish consumption habit (species and
 964 amount), age, gender, alcoholic consumption, health condition, economic level, etc.

965 **Biota**

- 966 24. Biota samples can provide information for different outcomes. Three types of outcomes,
 967 namely human exposures, environmental health, and temporal trends are identified in
 968 relation to biota monitoring. There is enough biotic mercury data available regionally and
 969 globally to assess environmental exposure for spatial and temporal trends for many, but
 970 not all, ecosystems and biomes of geographic interest. Human exposure to dietary
 971 methylmercury can originate from fish, birds and marine mammals (with fish forming a
 972 major contribution, birds forming either a minor or a major component, depending on
 973 diets, and marine mammals which can form a major contribution in certain diets).
- 974 25. The following samples from four major biomarker groups (taxa) are considered the most
 975 relevant and are most frequently used for methyl mercury monitoring:
- 976 ▪ Fish: muscle fillet, muscle biopsy, fin clips, blood
 - 977 ▪ Sea turtles: scutes, blood, muscle
 - 978 ▪ Birds: blood, feather, eggs, muscle, eggshells and membranes, liver and kidney

⁵ Assessment of prenatal exposure to mercury: human biomonitoring survey (2018) - the first survey protocol
<http://www.euro.who.int/en/health-topics/environment-and-health/chemical-safety/publications/2018/assessment-of-prenatal-exposure-to-mercury-human-biomonitoring-survey-2018>

979 ▪ Mammals: skin, fur or hair, muscle, liver and kidney
980 26. In assessing samples, it is recommended to assess muscle tissues for fish and marine
981 mammals. For birds, blood should be used for short term data, muscle or eggs should be
982 used for medium term and feathers can be used for long term results. It is considered to be
983 sufficient to assess total mercury for all tissues (assuming greater than 80 per cent
984 methylmercury mean level) using either wet weight or dry weight. Samples should be
985 georeferenced, with the level of detail varying according to the objective of the sampling.
986 Standard operating procedures are available for example through national /regional
987 monitoring programs, however additional more universal protocols may need to be agreed
988 on for other sampling which is not covered by this process. Inter-tissue conversions are
989 generally feasible to help provide a way to have standardized, and therefore comparable,
990 tissue mercury concentrations.

991 27. ↓
992 28. GBMS database was also the basis for the UN Environment’s Global Mercury Assessment
993 – 2018. Examples featured within the GBMS database include datasets for some
994 geographic areas with extensive temporal and spatial information, including areas for
995 freshwater lakes in the northern United States, much of Canada, and Scandinavia. These
996 areas represent over 500,000 fish mercury concentrations over the past 50 years of data
997 collection – sometimes with standard species. In order to potentially explain how the
998 temporal trends of fish mercury concentrations change under influence of different drivers,
999 including environmental/climate change in addition to deposition change, a set of
1000 minimum target information should be developed. For each location this should include
1001 lake (or river, estuary, sea etc.) catchment morphology, pollution deposition patterns, and
1002 local pollution history. For each biota species (here exemplified by fish) minimum data
1003 must include length, weight, sex, and sexual maturity. Samples (i.e. fish muscle) for
1004 determination of total mercury concentrations, may also be analysed for stable isotopes (at
1005 least nitrogen and potentially also carbon) for a better understanding of the food web
1006 processes. Many of these parameters are lacking from current databases. As an example,
1007 inter-annual and intra-annual variability is often much larger than long-term trends,
1008 making it difficult to relate temporal trend changes to large environmental drivers
1009 (including deposition). The spatial variation within the temporal trend must be considered
1010 when investigating convention effectiveness in years to come. To be able to document
1011 potential temporal trends changes, one need to lower the within-year variability, by
1012 improving the data adjustment, include more lake data and information, and collect data
1013 from the same lake over time.

1014 **Water and soil**

1015 29. Levels of mercury and mercury compounds in water are collected in relation to water
1016 quality issues in a number of countries. As the water release is considered as the another
1017 major part of anthropogenic emissions/releases together with atmospheric emission, those
1018 data is important in tracking mercury trends through appropriate data summary managing
1019 the impact of local activities. The mercury inflow to coastal area by rivers will influence
1020 the mercury methylation which may affect the people relying on the coastal subsistence
1021 fishing. Thus, although such impacts is local, the result will be used for global analysis on

Commented [m6]: Moving to INF suggested: It should be moved to the INF document as all other specific database/network information is now moved to INF.

Deleted: Biodiversity Research Institute (BRI) has compiled mercury data from published literature into a single database, the Global Biotic Mercury Synthesis (GBMS) Database. This database includes details about each organism sampled, its sampling location, and its basic ecological data. From each reference, mercury concentrations are averaged (using weighted arithmetic means) for each species at each location. Data have been compiled from 1,095 different references, representing 119 countries, 2,781 unique locations, and 458,840 mercury samples from 375,677 total individual organisms (See UNEP/MC/COP.3/INF/XX⁶).

Commented [m7]: Revision suggested: The description does not properly address the importance and usefulness of the water data. Research based programmes also provide important and useful information. Non-existence of long-term monitoring programmes is not the reason to disqualify the usefulness of the information.

vulnerable population as the coastal area provides subsistence fishery (in comparison with pelagic fishery).

30. Levels of mercury in ocean water could be comparable on a global basis and collected by existing networks and ad hoc research programmes. Understanding mercury transfer from mercury emission to air, then to ocean, fish and finally to human is essential to evaluate mercury risks.

31. Soil samples may be very useful in assessing the state of contamination of a particular site, but global comparability may not be feasible, given differences in soil types etc. Data on the levels of mercury in sediments are very relevant for the associated levels of mercury in biota; however sampling of sediment was considered not as widespread, nor as easily comparable on a global basis, at this time.

Deleted: These data may be useful in tracking mercury resulting from local activities which release mercury; however, will not provide overall trends on a global

Deleted: , but currently such work is done through research-based activities and not dedicated long term monitoring programmes. ...

Deleted: C

Deleted: urrently, this work is done through research-based activities and not dedicated long term monitoring programmes....

3. Comparability, gaps and options for filling gaps

Air

32. Figure 1 shows the current monitoring efforts for TGM/GEM. From this figure it can be seen that the gaps of TGM/GEM information could be filled by enhancing the current networks that are conducting atmospheric mercury monitoring. Such expansions would include areas within South America, Africa, the Caribbean, parts of Asia, Russia, and Oceania.




Figure 1 – Existing monitoring networks measuring Hg concentrations in air.

33. The following are recommended:

- Couple current monitoring of TGM/GEM with new technologies (including passive and active mercury sampling);
- Expand current monitoring networks, where possible, to fill in data gaps;

- 1087 ▪ Employ currently-used standard procedures for data collection and treatments, where
- 1088 possible;
- 1089 ▪ Conduct intercomparisons of measurement technologies and data treatment among
- 1090 networks;
- 1091 ▪ Fill geographical data gaps of information using manual active or passive sampling
- 1092 methods;
- 1093 ▪ If feasible, couple manual active or passive air measurements with active and wet/dry
- 1094 deposition measurements;
- 1095 ▪ Conduct sampling at least on a quarterly basis (either averaged with active sampling
- 1096 data or integrate over 3 months with passive sampling) to assess seasonal variation;
- 1097 ▪ Prioritize gaps identified in the global mercury assessment and other literature for the
- 1098 establishment of new site locations.

1099 33. In elaborating future strategies aiming to fill geographical gaps of atmospheric mercury
1100 monitoring data it is recommended to ensure the operation of about 30 monitoring sites
1101 with manual active or passive air sampling in large geographical areas such as Africa,
1102 Latin America and Russia placed in locations that may provide information on regional /
1103 local background Hg concentrations. The suggested number of sites is only indicative: a
1104 larger number of sites using manual active or passive air sampling in these areas would
1105 certainly allow to have a better geographical distribution and representativeness of the
1106 regional/local emission regimes, meteorology and transport/deposition patterns. A cost
1107 analysis for air monitoring including the proposed sampling can be found in
1108 UNEP/MC/COP.3/INF/XX Part I Section 4.

1109 **Humans**

1110 34. Studies using the WHO protocol for assessment of prenatal exposure to methylmercury
1111 are recommended to fill the data gaps in order to obtain a global picture necessary for
1112 effectiveness evaluation. The protocol enables collection of comparable data (e.g. hair
1113 samples from 250 people per study location with minimum diversity recommended). The
1114 studies are country-driven. Local ethical (Institutional Review Board) clearance is
1115 required and the studies are conducted within the health system, therefore country
1116 approval is a given. Each country owns its data and submission of results is voluntary.

1117 35. Article 17 of the Minamata Convention on Mercury specifies in paragraph 1(d) that each
1118 party shall facilitate the exchange of epidemiological information concerning health
1119 impacts associated with exposure to mercury and mercury compounds, in close
1120 cooperation with the World Health Organization and other relevant organizations, as
1121 appropriate. The compilation and exchange of data on mercury levels obtained through
1122 human biomonitoring should be undertaken in line with this article of the Convention.

1123 36. To facilitate the generation of globally representative data and trend information on human
1124 biomonitoring, which will be most relevant for effectiveness evaluation, an oversight body
1125 should be kept informed of the studies planned and carried out.

1126 37. Data quality issues are covered by the WHO protocol. Results of the measurements must
1127 be analytically comparable between laboratories/different studies. To ensure
1128 comparability, each national survey would need to follow the WHO harmonized SOPs for
1129 sampling and analytical methods, and develop procedures for quality assurance and
1130 quality control that cover the pre-analytical phase. The availability of appropriate

1131 reference materials (samples with a certain level of mercury)⁷ supports internal quality
1132 assurance. External quality assurance should be done through international inter-
1133 laboratory comparison investigations. Coordination of the studies will contribute to ensure
1134 appropriate quality control measures.

1135 38. The WHO protocol also covers data management, analysis and evaluation issues,
1136 including whether this should be done at the national and/or international level. It
1137 recommends that participating countries conduct statistical analyses at the national level
1138 and submit anonymized data for statistical analysis to a central database. The aim of a
1139 statistical analysis at the international level is to assess associations between biomarker
1140 values and predictors such as age, gender, fish consumption habits, etc. (collected via
1141 questionnaire) in a pooled dataset. Data communication issues are also addressed in the
1142 WHO protocol and particularly for indigenous peoples in AMAP Human Health
1143 Assessments. These communication issues include communication of the results within
1144 the country, to the individuals participating in the study and to policy makers. It should be
1145 noted that, in some countries, national guidelines relating to communication of results may
1146 already exist.

1147 39. ↓
1148 **Biota**

1149 40. It has been recognized that there is a large amount of published data available, as well as
1150 unpublished data collected for commercial and governmental purposes. However, it is not
1151 clear to what extent published and other data reflect background information on mercury
1152 concentrations, or whether existing data emphasizes areas where high mercury
1153 concentrations are expected. As previously described, the large , biotic mercury
1154 concentration datasets from the northern United States, Canada and Scandinavia revealed
1155 that levels in freshwater fish from lakes with local mercury sources responded to
1156 regulation and management. Further evaluation work on existing data is required to gather
1157 all currently available globally representative biotic mercury data, to assess what data are
1158 relevant, comparable and able to be harmonized. This process has been started with the
1159 partly UNEP funded GBMS dataset, which will allow a clearer identification of data gaps,
1160 which may be geographic or taxonomic.

1161 41. The Arctic Monitoring and Assessment Programme (AMAP) is one of the best examples
1162 of how to operate a long-term Hg biomonitoring field program for the benefit of both
1163 human and ecological health (AMAP 2011, 2015). Whereas, the WHO Global
1164 Environment Monitoring System - Food Contamination Monitoring and Assessment
1165 Programme, commonly known as GEMS/Food, has one of the best global systems for
1166 collecting fish Hg data through their network of collaborating centers and recognized
1167 national institutions (WHO 2018).

1168 **Water**

1169 42. Global data in rivers flowing into ocean is especially important as there is substantial
1170 uncertainty in estimates of global riverine discharge as compared with the atmospheric
1171 deposition which are well established. The mercury data in water are usually collected in

Deleted: The UNEP/WHO GEF Global Monitoring Project demonstrated generation of data using the WHO Protocol in developing countries to be cost-effective, practical and feasible. The project built local capacities to conduct such studies, which can therefore be repeated over time and in a range of locations to fill gaps, as described in paragraph 20. ...

Commented [m10]: New paragraphs suggested: Comparing previous 3 media, i.e. air, biota and human, larger gap exists which should be described in this section.

⁷ A list of existing reference materials can be found in UNEP/MC/COP.3/INF/XX Part II.

1179 relation to local water quality and thus not globally harmonized and will not provide
1180 overall trends on a global basis.
1181 43. Understanding mercury transfer from mercury emission to air, then to ocean, fish and
1182 human is essential to evaluate mercury risk. Among them, the largest gap exists for marine
1183 data. Levels of mercury in ocean are collected by existing networks and ad hoc research
1184 programmes, but currently such work is done through research-based activities and not
1185 dedicated long term monitoring programmes. As current data gap is the highest for marine
1186 monitoring, it will contribute the most for improving the precision of the overall results.

1187 **Cost analysis**

1188 44. A table summarizing the cost, practicality, feasibility, sustainability, comparability and
1189 coverage of currently-used monitoring methods for air, humans, biota and water is
1190 included in UNEP/MC/COP.3/INF/XX.

1191

1192 **4. Available modelling capabilities to assess changes in global mercury levels within and** 1193 **across different media**

1194 45. Table 1 summarizes the capabilities of models to assess changes in global mercury levels
1195 within and across different media. Models for different media (air, water, land, biota) vary
1196 in their ability and state of development. Atmospheric models have been extensively
1197 evaluated and can be applied to assess spatial gradients in atmospheric mercury
1198 concentrations and deposition, as well as temporal changes. By contrast, models for other
1199 media such as land are still mainly used in research applications. Further explanation
1200 including reference to specific available models and example geographic presentation of
1201 calculations from existing models can be found in UNEP/MC/COP.3/INF/XX.

1202 46. Integrated modeling frameworks can illustrate pathways by which primary releases of
1203 mercury to the atmosphere, land and water reach methylmercury in fish and wildlife as
1204 well as exposure of some fish consuming human populations. At present, integrated
1205 modeling frameworks are under development and available as a research product.
1206 Integrated models have not previously been applied or compared in global assessment
1207 efforts. Coupled atmosphere-ocean and atmosphere-terrestrial have been published in the
1208 peer-reviewed literature by a few research groups. With additional model evaluation,
1209 updates should be available to begin policy-relevant analyses by 2023. Models for food
1210 web bioaccumulation of methylmercury are also available from selected groups and can be
1211 used to describe accumulation patterns at the ecosystem scale (lakes, wetlands, estuaries,
1212 contaminated sites) and for global marine food webs. The most difficult link in integrated
1213 modeling frameworks is to human exposure and health outcomes due to the diversity of
1214 dietary preferences, food consumption patterns and individual variability in toxicokinetics
1215 affecting methylmercury uptake and elimination. All these components of integrated
1216 modeling frameworks are rapidly developing in the scientific community.

1217

1218

1219
1220

Table 1. Summary of available modeling capabilities for individual media.

Media/Availability	Indicators needed for model input	Output provided	Gaps still to be filled
<p>Socio-Economic Modeling: Some Availability</p> <p>Global emission models (forecasting up to 2050)</p>	<p>Inputs: socio-economic activity data (production, population, GDP), material flow and policy specifications</p> <p>Evaluation: intercomparison and past performance, Anthropogenic material flow</p>	<p>Global demands, Emission and release scenarios</p>	<p>Mercury emission factors to be refined(regional, site, etc.), data on mercury content of commodities to be collected, consistency across sectors and non-mercury policies to be explored (e.g. energy)</p>
<p>Air: Widely Available</p>	<p>Inputs: Global emissions</p> <p>Evaluation: Atmospheric measurements; Wet and dry deposition data</p>	<p>Atmospheric concentration; deposition; temporal changes; attribution by source region</p>	<p>Harmonized emissions inventories to be established</p>
<p>Water: Research Product; Some Availability</p> <p><i>Global Oceans</i> Global ocean models (MITgcm, NEMO model)</p> <p><i>Estuaries (site specific); Freshwater/rivers (site specific)</i></p>	<p>Inputs: Spatially resolved global atmospheric Hg inputs (wet + dry) – Concentrations of Hg and MeHg in rivers (globally)</p> <p>Evaluation: Measured seawater total and methylmercury, and Hg⁰ concentrations; These are being collected through networks (GEOTRACES/CLIVAR)</p>	<p>Seawater MeHg in global oceans</p> <p>Total Hg concentrations in seawater globally for surface/deep ocean</p> <p>Temporal changes</p>	<p>Seawater Hg species data somewhat sparse but improving</p> <p>Data on Hg and MeHg in global rivers largely lacking</p>
<p>Soils/Land: Research Product – Some Availability</p> <p><i>Global soils</i> Global terrestrial mercury model (GTMM)</p>	<p>Inputs: Atmospheric deposition (model input) Emissions releases to land/water (very preliminary and coarse spatial resolution)</p> <p>Few data on runoff from contaminated sites</p>	<p>Soil Hg concentrations globally</p> <p>Hg in global rivers</p>	<p>MeHg simulation for terrestrial environments other than site specific assessments still to be done.</p>

ASGM/Contaminated sites (not yet integrated into global models but would be useful)	Global land cover data and atmospheric inputs Evaluation: Soil Hg data (good data for North America, parts of Europe)	“Hot spots” most sensitive to Hg inputs and likely to affect biota/human populations	Ground truthing global “hot spot” analysis is needed. Data on locations of ASGM and releases/contaminated sites to be collected.
Biota: Research Product – Some Availability Food web bioaccumulation model for marine ecosystems (global models for plankton exist/fish under development) Food web model for freshwater ecosystems (site specific)	Inputs: MeHg seawater (model); Fisheries biomass production from primary productivity globally, trophic interactions (available through collaboration with NOAA/GFDL and UBC Nereus projects) Evaluation: Biotic mercury database Trophic level 3 for temporal trend, 4 for spatial gradient analysis	Concentrations of MeHg in fish consumed by human populations; Marine origin of MeHg and Attribution of Hg sources in fish (marine mammals?) by region; Changes due to emissions and climate	Global fish model under development; could link to marine mammals/birds Trophic level 4 data in Asia and Africa to be collected
Humans Exposures of marine fish consumers (globally) <i>Toxicokinetic model linking MeHg ingestion and blood/hair concentrations/outcomes</i> Freshwater fish and rice consumers (site specific data, if applicable) – these may be highest risk populations	Inputs: Biomass and MeHg concentrations in fish consumed by different subsistence populations globally (model); Dietary intake data for different human populations National biomonitoring data (model evaluation)	Goal: Attribution of Hg source contributions to human populations	Mechanisms affecting relationships between external MeHg exposure and blood concentrations/outcomes for different populations are uncertain (research evolving)

Occupational exposures at ASGM sites (site specific)			
--	--	--	--

1221
1222

1223 **5. Establishing a baseline for monitoring data**

1224 45. In the “before-after” approach where the mercury levels before and after the
1225 implementation of the Convention, monitoring data close to the beginning and the end of
1226 the evaluation period can be used. For the first effectiveness evaluation, monitoring data
1227 before the entry into force of the Convention may be used as baseline.

1228 46. For air, historical monitoring data exist for some part of northern hemisphere. For human
1229 biomonitoring, data from a limited number of regional and national biomonitoring
1230 programmes and longitudinal studies may be used. For biota, historical data on mercury
1231 levels in freshwater fish in limited geographical areas are available. Work is underway to
1232 analyze available data on ocean fish species.

1233 47. In the “with-without” approach to assess the change in mercury levels attributable to the
1234 measures taken to implement the Convention, mercury levels for the business-as-usual
1235 scenario need to be estimated using integrated modelling framework described above.

1236

1237 **Annex 2: Draft terms of reference of the Effectiveness Evaluation Committee**

1238

1239 **A. Mandate**

1240 1. An Effectiveness Evaluation Committee (hereinafter, “the Committee”) is
1241 established to perform the functions assigned to it by the Conference of the Parties.

1242 **B. Membership**

1243 2. The members of the Committee shall be appointed on the basis of equitable
1244 geographical distribution, taking into account gender and the need for a balance
1245 between types of expertise.

1246 3. The Committee shall consist of twelve experts, as follows:

1247 (a) Ten experts designated by parties representing the five United Nations
1248 regions, and confirmed by the Conference of the Parties;

1249 (b) One expert representing the monitoring arrangement;

1250 (c) One expert representing the implementation and compliance committee.

1251 4. Experts designated by parties and confirmed by the Conference of the Parties
1252 shall have expertise in evaluation, reporting and national implementation, financial
1253 or technical assistance, or other expertise relevant to the evaluation.

1254 5. Experts from the implementation and compliance committee shall be selected
1255 by and from among the members of its committee.

1256 5a The expert representing the monitoring arrangement shall be selected from the
1257 members that take part in these arrangements.

1258 5b Members shall provide their expertise in a neutral and impartial manner, and
1259 stand to the evidence presented to the committee.

1260 6. The terms of office shall coincide with a cycle of evaluation as determined by
1261 the Conference of the Parties.

1262 7. If a member is unable to complete his or her term of office, the region
1263 nominating that member shall nominate another person to complete the term.

1264 **C. Invited experts and observers**

1265 8. The Secretariat shall select two internationally recognized experts in
1266 effectiveness evaluation with due consideration to available expertise on the
1267 measures.

1268 9. The Secretariat shall invite one representative of the World Health
1269 Organization as an observer.

1270 10. The committee will invite the participation of up to five experts from civil
1271 society, indigenous organizations, intergovernmental organizations, industry and the
1272 UNEP Global Mercury Partnership as observers. The participation of observers will
1273 be balanced among the above-mentioned groups and gender.

1274 11. The committee may allow additional observers within reasonable limits.

1275 12. Observers shall provide their technical expertise that helps the committee
1276 members interpret the information provided.

1277 **D. Officers**

1278 13. The committee shall elect, from among its members, a chair and a vice-chair.

1279 **E. Administrative and procedural matters**

1280 14. The committee shall apply, mutatis mutandis, the rules of procedure of the
1281 Conference of the Parties, unless otherwise provided in these terms of reference.

1282 15. The committee may establish such arrangements as are necessary to facilitate
1283 its work in line with the present terms of reference.

1284 16. The committee members shall seek to reach agreement by consensus. Should
1285 consensus not be reached by members, the range of their views shall be reflected in
1286 any report to be submitted to the Conference of the Parties.

1287 **F. Meetings**

1288 17. The committee shall hold two face-to-face meeting, to review the information
1289 available for each evaluation cycle and to develop a report to the Conference of the
1290 Parties, subject to the availability of funds and work requirements. Based on the
1291 decisions of the Conference of the Parties, the frequency of committee meetings
1292 may be amended as necessary.

1293 18. Documents to be transmitted to the Conference of the Parties shall be
1294 finalized by the committee at least four months before the meeting of the
1295 Conference of the Parties.

1296 **G. Language of meetings**

1297 19. The working language of the committee shall be English.

1298 **H. Budget**

1299 20. Except for members from developed country parties referred to in paragraph 4
1300 of the present terms of reference, financial support for travel and daily subsistence
1301 allowance shall be made available to committee members, and invited experts and
1302 observers for participation in meetings of the committee according to United
1303 Nations rules and practice.

1304

1305 **Annex 3: Draft terms of reference of the global monitoring arrangements**

1306

1307 **Introduction**

- 1308 1. This annex contains a proposal for global monitoring arrangements building on existing
1309 monitoring activities, knowledge, expertise and proposes the terms of reference for an
1310 expert group to [prepare a synthesis report on monitoring as identified in Section III of
1311 the report] [carry out tasks related to monitoring indicators identified in the effectiveness
1312 evaluation framework in Section III] in this Annex.
- 1313 2. In the consideration of the monitoring arrangements, the following key elements were
1314 identified:
- 1315 a. Mercury data and their availability from human health and environmental
1316 monitoring programmes that achieve global coverage and contain at least core
1317 representative data from all regions,
- 1318 b. Tools supporting data harmonization such as standard operating procedures and
1319 monitoring guidance document,
- 1320 c. Expertise necessary for gathering and consolidating harmonized information that
1321 ensures comparability and consistency in mercury monitoring data over the long-
1322 term,
- 1323 d. Modelling capabilities, and
- 1324 e. Development of a global periodic report on levels and trends of mercury to support
1325 the effectiveness evaluation.
- 1326 3. The text below further elaborates on the key elements identified, but a large amount of
1327 other relevant technical information on monitoring and background complementing the
1328 proposal below is available in a reference document as UNEP/MC/COP.3/INF.xx.
1329 Existing modelling capabilities are reviewed in detail in that INF document as well.

1330 **1. Mercury data and their availability from human health and environmental monitoring**
1331 **programmes**

- 1332 4. Regarding mercury data availability, a review presented in Annex I shows that even if
1333 mercury has one of the largest available collective data sets of recognized environmental
1334 contaminants, data gaps remain. These gaps could be efficiently covered with support of
1335 scientific activities and use of already developed materials.
- 1336 5. By continuing existing mercury monitoring activities in a harmonized manner (see Tools
1337 supporting data harmonization below), supplementing them with actions to fill the
1338 geographical gaps, data on levels of mercury and mercury compounds in **environment,**
1339 **biotic media and vulnerable populations** either are available or would be able to be
1340 obtained, and would be comparable on a global basis.

Commented [m11]: Word change suggested: To be consistent with the wording used in the Article 22 of the Convention.

Deleted: air, biota and humans

1342 6. Below is the proposal for mercury monitoring activities building on existing monitoring
1343 activities and knowledge organized by media. This information should be part of global
1344 mercury monitoring report.

1345 **Air**

1346 7. For air monitoring, it is proposed to continue monitoring activities by existing networks
1347 by active continuous monitoring and manual active and passive air sampling techniques
1348 and collect:

- 1349 ▪ Total Gaseous Mercury (TGM) concentrations in air at background and
1350 impacted sites, and
- 1351 ▪ Atmospheric deposition fluxes.

1352 for assessing spatial and temporal patterns of mercury concentrations in ambient air and
1353 deposition fluxes to terrestrial and aquatic ecosystems. Standard Operating Procedures
1354 (SOPs) suggest to monitor mercury deposition fluxes with samplers that are “Wet only”
1355 or Bulk.

1356 8. To fill the geographical gaps in continents, samples should be collected to provide
1357 average TGM concentrations as monthly (or at lower frequency) average to cover Africa,
1358 Latin America and Russia and provide regional/local background TGM concentrations.

1359 **Human**

1360 9. For human biomonitoring for prenatal exposure in the general population the following
1361 biomarkers are recommended:

- 1362 ▪ Total mercury in maternal scalp hair (3 cm hair strand from the scalp, to
1363 measure exposure during the 3rd trimester), and
- 1364 ▪ Total mercury in cord blood – recent exposure to methyl mercury.

1365 10. Maternal scalp hair is a preferable biological matrix to assess prenatal exposure. Cord
1366 blood can be an alternative matrix to hair. Human samples collected in approximately 5-
1367 year intervals are feasible for human biomonitoring surveys considering the aim to
1368 identify statistically significant differences as well as the time such studies take to
1369 implement (including adaptation of the master protocol to local circumstances, local
1370 ethical approval, training of staff etc.). Human samples should be accompanied by a
1371 series of attributes, e.g. age, gender as well as social/habitual information e.g. fish
1372 consumption pattern, economic level, etc.

1373 11. It might be useful to coordinate the sample collection with the survey activities under the
1374 Stockholm Convention as the one ethical approval could be used.

1375 12. The Global Mercury Assessment 2018 identified currently available data on mercury
1376 exposure in national human biomonitoring programmes, longitudinal birth cohort studies
1377 and cross-sectional information in specific populations including high exposure groups.
1378 These activities should be continued to provide a long-term information for subsequent
1379 effectiveness evaluation.

1380

Biota

1381 13. For biota monitoring, an important aspect in combining monitoring efforts for
 1382 documentation of convention effectiveness would be to define biological species and
 1383 proper tissue types for monitoring, to minimize the effects of species-specific
 1384 physiological differences. Species that accumulate significant amounts of mercury pose a
 1385 potential risk for human health, that are widely distributed over specific geographically
 1386 areas, and that exist in numerous historical studies should be prioritized. Additionally,
 1387 there is a need to normalize or account for mercury concentrations in biota by size, age
 1388 and sex, and these data should be included in the data collection process. The choice of
 1389 fish species for sampling should be based on the purpose of the sample use. Human
 1390 biomonitoring should be accompanied by the fish consumption pattern, thus, mercury
 1391 levels in typical commercial fishing species are appropriate, related to human and
 1392 ecological health assessments.

1393 14. It is proposed that biotic monitoring be separated into two major approaches to account
 1394 for major differences in exposure pathways: continental and oceanic frameworks. A large
 1395 amount of relevant technical information on the frameworks is available in a reference
 1396 document as UNEP/MC/COP.3/INF.xx. Continental framework aims at identifying
 1397 ecosystem sensitivity spots that are able to methylate mercury and make it available in
 1398 the food web. Oceanic Framework for mercury monitoring in biota covers oceanic areas.
 1399 The outcome combines ocean basin, matrix of interest for human consumption that have
 1400 global ranges to define spatial gradients (trends) of mercury level in biota.

1401

Water

1402 15. There are GEOTRACES and CLIVAR programs, and ad hoc research programs for
 1403 marine monitoring. While development of an enhanced database on speciated mercury
 1404 concentrations in seawater is strongly encouraged, such measurements are typically
 1405 collected by analytical specialists to ensure data quality.

1406

2. Tools supporting data harmonisation

1408 16. Tools supporting data harmonization regarding comparability represent in particular
 1409 standard operating procedures, guidance on global monitoring document, and inter-
 1410 calibration studies.

1411 17. Document UNEP/MC/COP.3/INF/XX contains a more detailed information on standard
 1412 operation procedures (SOPs) already available and their use is encouraged. Review of
 1413 data availability therein also comprises information on other available tools for
 1414 maintaining data comparability including inter-calibration studies.

1415 18. Further, to maintain harmonized information on mercury levels in environment, existence
 1416 of a global mercury monitoring guidance document would be very useful. While
 1417 development of such a document is included in the ad-hoc expert group`s mandate, the
 1418 group felt that such document can only be prepared once monitoring arrangements for
 1419 mercury are agreed. Guidance document could then be prepared swiftly on the basis of
 1420 core matrices and available knowledge.

Commented [m12]: Revision suggested: For assessing human exposures, the mercury levels of commercial fish are important.

Deleted: the trophic level, with trophic level 4 (carnivores that eat other carnivores) being most appropriate for decisions ...

Commented [m13]: New paragraph suggested: Some ongoing programmes actually collecting mercury data in water.

1424 19. Nevertheless, experts prepared elements for the guidance on global monitoring (available
1425 monitoring activities organized per matrix, state of science for monitoring, procedures on
1426 sampling, sample handling, chemical analyses of samples) that is contained in
1427 UNEP/MC/COP.3/INF/XX part two that presents a draft structure of the guidance
1428 document and other relevant information.

1429

1430 **3. Development of a global periodic report**

1431 20. It is proposed that a global mercury monitoring report on status of the environment and
1432 occurrence of mercury is developed in regular and suitable intervals to support the
1433 effectiveness evaluation.

1434 21. Available globally representative monitoring data would be compiled, assessed and
1435 summarized by relevant experts performing scientific function in this field (see below).

1436 22. Global report would be organized by media and show available monitoring data and
1437 trends in the environment, humans and biota. Global monitoring report would also use
1438 models to predict further trend development.

1439 23. Information from the global report would then be also used for contextualization of
1440 information in a multi compartment model to capture the socio-economic scenario,
1441 baseline and different policy alternatives.

1442 24. The first global report on monitoring and modelling to the effectiveness evaluation
1443 committee on state of the environment needs to become available for the first meeting of
1444 the effectiveness evaluation committee.

1445

1446 **4. Expertise necessary for synthesizing monitoring data**

1447 25. During the discussions of the effectiveness evaluation framework`s science and technical
1448 functions, it became clear that information on the status of the environment and
1449 occurrence of mercury is to be synthesized by an expert body with extended research
1450 expertise to oversee the gathering and consolidation of monitoring data.

1451 26. The group would be assigned to gather information from existing monitoring activities
1452 and compile them into a global synthesis report and assess mercury levels and trends
1453 through the use of models, and thus prepare a global monitoring report as referred to in
1454 Section II of this report. Proposed terms of reference of the group are shown below.

1455

1456 **Draft ToR of the monitoring and modelling group**

1457 **Mandate**

1458 A monitoring and modelling [task] group (hereinafter, “the group”) is established to perform
1459 the functions assigned to it by the Conference of the Parties in support of the effectiveness
1460 evaluation including:

- 1461 a. Gathering of information from mercury monitoring activities and compilation of the
1462 relevant information including national and scientific data on changes in levels of
1463 mercury in core media taking into account the work already achieved and drawing on
1464 experience from existing monitoring networks on mercury. Changes include spatial and
1465 temporal trends including contextualization through use of models.
- 1466 b. Preparation of a global monitoring report on mercury for effectiveness evaluation
1467 committee meetings.
- 1468 c. Development of a monitoring guidance document to provide the COP with
1469 comparable monitoring data on the presence and movements of mercury and mercury
1470 compounds in the environment as well as trends in levels of mercury and mercury
1471 compounds observed in biotic media and vulnerable populations, organize data gathering
1472 and visualization of information. The group should start its work on this task immediately
1473 so that the documents is available for COP4.
- 1474 d. Update of a monitoring guidance document in line with the latest scientific
1475 knowledge, modelling capabilities and ongoing monitoring activities.
- 1476 e. Identification of gaps in information/knowledge and development of proposals for
1477 bridging the gaps as a part of the report prepared for consideration by the effectiveness
1478 evaluation committee.

1479 **Membership**

1480 The group members shall be appointed on the basis of equitable geographical distribution,
1481 taking into account gender and the need for a balance between types of expertise.

1482 Each region should nominate four experts for monitoring and modelling (up to three
1483 representatives with expertise on mercury monitoring in core media or participating in
1484 existing monitoring networks on mercury and at least one representative with expertise on
1485 modelling environmental trends/multicompartment models)

1486 The group will invite the participation of up to 10 experts from civil society, indigenous
1487 communities, intergovernmental organizations, industry and “global modelling” experts. The
1488 participation of these experts as observers will be balanced among the above-mentioned
1489 groups.

1490 The group will invite relevant experts from research communities, Global Mercury
1491 Partnership and existing monitoring networks to assist them in their work and supplement the
1492 most up to date information and scientific knowledge to produce a global report.

1493 The terms of office shall coincide with a cycle of the effectiveness evaluation as determined
1494 by the Conference of the Parties. To maintain continuity, the COP may renew terms of office
1495 of the members for subsequent evaluations. If a member is unable to complete his or her term
1496 of office, the region/sector nominating that member shall nominate another person to
1497 complete the term.

1498 **Officers**

1499 Two co-chairs will be elected by the group to facilitate its meetings.

1500 **Secretariat**

1501 The secretariat will provide administrative and programmatic support to the group of experts.

1502 **Meetings**

1503 The group on monitoring and modelling will meet face-to-face at least three times during an
1504 effectiveness evaluation cycle to coordinate monitoring activities on mercury and to deliver a
1505 global report on monitoring and modelling to the effectiveness evaluation committee on state
1506 of the environment.

1507 **Language**

1508 English will be the working language of the group.

1509 **Annex 4: Description of the scientific and technical functions**

1510
1511

- 1512 1. The framework foresees five reports being prepared for consideration by the
1513 Effectiveness Evaluation Committee (see Section III of the Report).

1514 **Synthesis Reports**

- 1515 2. Four synthesis reports are to be prepared (level 1 to 3) to respond to the first three policy
1516 questions. Two streams of information feed into these reports: (i) information provided
1517 by Parties based on Article 21 reporting, and (ii) information and knowledge that is
1518 scientific, peer-reviewed and publicly available.

- 1519 3. The content envisaged under each synthesis report is set out below, as are the tasks that
1520 need to be completed, and the expertise required.

1521 i. The **Emissions and Releases Report** is to gather, analyse and synthesise relevant
1522 information on emissions and releases inventories from relevant sources, as
1523 specified in Article 8 and 9, as well as information on the measures taken by
1524 Parties to control mercury emissions and releases, and relevant changes in
1525 emissions and releases. The expertise required for this task includes
1526 emissions/releases inventories, developing or implementing measures to control
1527 mercury emissions and releases from relevant sources, including best available
1528 techniques and best environmental practices, modelling and inventories on
1529 temporal and spatial trends and variability.

1530 ii. The **Trade, Supply and Demand Report** is to gather, analyse and synthesise
1531 relevant information on the mercury flows and social stocks, on trends in trade,
1532 supply and demand for mercury, and on regulatory frameworks and
1533 implementation. The expertise required for this task includes: trade analytics,
1534 sectoral analysis, ASGM expertise, use, changes and alternatives to mercury in
1535 products and processes.

1536 iii. The **Waste Management Report** is to gather, analyse and synthesise relevant
1537 information on mercury waste flows and stocks, track mercury waste management
1538 practices and recycling, and on regulatory frameworks and implementation, as
1539 well as gaps. The expertise required for this task include: inter-industry relation
1540 analysis, waste management policy and practices, and waste disposal engineering.

1541 iv. The **Global Monitoring Report** is to gather information from mercury monitoring
1542 activities and compile relevant information including national and scientific data
1543 on changes in levels of mercury, taking into account the work already achieved,
1544 and drawing on the experience of existing networks on mercury, trends and
1545 models. (See Annex III for detailed information).

1546 **Integrated Assessment Report**

- 1547 4. Based on the synthesis reports and other information linkages need to be made between
1548 policy actions, emission reductions and resulting mercury levels, using available data

1549 sources, modelling techniques and analytical tools drawn from natural and social
1550 sciences. This will be done in the **Integrated Assessment Report**.

1551 5. It is to be noted that the integrated assessment function will evolve as our understanding
1552 of mercury improves over time. For the first round of the effectiveness evaluation, when
1553 no previous assessment is available, several ground studies to provide the basis of the
1554 evaluation, will need to be conducted.

1555 6. With this as background, the content of the integrated assessment report is expected to
1556 contain:

1557 a. The examination of time lags between actions and outcomes observed by the
1558 subsequent evaluations: Significant time lags for years or even decades due to the
1559 slow pace of change in socio-economic systems, and in the physical and
1560 biological dynamics of the Earth system will need to be discussed.

1561 b. The examination of the baseline scenario, which draws on a hypothetical
1562 'business as usual' setting for when the Convention had not been implemented:
1563 The hypothesis will employ assumptions and interpretations that could go beyond
1564 the factual presentation. As far as practical, different scenarios will be developed
1565 for future forecasting, given that it is expected that population growth, economic
1566 development, and global warming will alter the mercury baseline due to the
1567 changes in consumption patterns and global material flows.

1568 c. The assessment of the four policy questions, that could go as far as forecasting
1569 based on appropriate extrapolation: Several types of modelling can help such an
1570 assessment. (It has to be noted, however, that present science has not yet
1571 developed reliable models to forecast long-term changes in mercury levels
1572 resulting from emissions reductions that take into account the full complexities of
1573 mercury in the environment.) Therefore, earlier evaluations on the effectiveness
1574 of the Minamata Convention must rely on simpler forecasting methods and will
1575 have greater uncertainty than later evaluations when improvements to such
1576 forecasting models become available.

1577 d. The comprehensive analysis of the interaction between different indicators for
1578 identifying important synergies and trade-offs: Understanding the relationship
1579 between indicators is important for improving implementation efficiency.

1580 7. The following chapters are suggested for the Integrated Assessment Report:

1581 **Assumptions and baseline scenario setting for the integrated assessment**
1582 **Assessment of the policy questions**
1583 *Policy question 1:* Have the Parties taken actions to implement the Minamata
1584 Convention?
1585 *Policy question 2:* Have these actions resulted in changes in emissions and releases of
1586 mercury to the environment?
1587 *Policy question 3:* Have these changes in emissions and releases resulted in changes in
1588 levels of mercury in the environment, biota and humans attributable to the Convention?
1589 *Policy question 4:* Will existing measures under the Minamata Convention be sufficient
1590 to meet its objectives of protecting human health and environment from mercury?
1591 **Synergies and trade-offs between indicators for improving implementation**
1592 **efficiency**

1593 **Time lags between actions and outcomes**
1594 **Conclusions**
1595 **Appendix: Result “Dashboard” - progress of the indicators in the evaluation**
1596 **framework**
1597